



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/82, 15/29, C07K 14/415, C12N 5/10, C07K 16/16, C12Q 1/68, G01N 33/563, A01H 5/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/06754</b>  <b>(43) International Publication Date:</b> 10 February 2000 (10.02.00)
<b>(21) International Application Number:</b> PCT/NL99/00491  <b>(22) International Filing Date:</b> 30 July 1999 (30.07.99)  <b>(30) Priority Data:</b> PCT/NL98/00445 31 July 1998 (31.07.98) NL  <b>(71) Applicants (for all designated States except US):</b> CENTRUM VOOR PLANTENVEREDELINGS-EN REPRODUC-TIEONDERZOEK (CPRO-DLO) [NL/NL]; P.O. Box 16, NL-6700 AA Wageningen (NL). LANDBOUWUNI-VERSITEIT WAGENINGEN [NL/NL]; P.O. Box 8123, NL-6700 ES Wageningen (NL).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> VAN DER VOSSEN, Edwin, Andries, Gerard [NL/NL]; Bouwstraat 1, NL-3572 SN Utrecht (NL). VAN DER VOORT, Jeroen, Nicolaas, Albert, Maria, Rouppe [NL/NL]; Omval 91, NL-1096 AA Amsterdam (NL). LANKHORST, Rene, Marcel, Klein [NL/NL]; Nijburgsestraat 43, NL-6668 AZ Randwijk (NL). BAKKER, Jaap [NL/NL]; Geertjesweg 122, NL-6704 PD Wageningen (NL). STIEKEMA, Wilhelmus, Johannes [NL/NL]; Leonard Roggeveenstraat 21, NL-6708 SL Wageningen (NL).		<b>(74) Agent:</b> DE BRUIJN, Leendert, C.; Nederlandsch Octrooibu-reau, Scheveningseweg 82, P.O. Box 29720, NL-2502 LS The Hague (NL).  <b>(81) Designated States:</b> JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> ENGINEERING NEMATODE RESISTANCE IN SOLANACAE		
<b>(57) Abstract</b>  <p>The present invention relates to the <i>Gpa2</i> resistance gene from potato conferring resistance to phytopathogenic nematodes of the genus <i>Globodera</i>. It further relates to methods and materials employing the gene and processes for identifying related genes. Finally the invention relates to polypeptides encoded by said resistance genes and the use of said polypeptides.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

## ENGINEERING NEMATODE RESISTANCE IN SOLANACAE

### FIELD OF THE INVENTION

5 The present invention relates to the *Gpa2* resistance gene from potato conferring resistance to phytopathogenic nematodes of the genus *Globodera*. It further relates to methods and materials employing the gene and processes for identifying related genes. Finally the invention relates to polypeptides encoded by said resistance genes and the use of said polypeptides.

### BACKGROUND OF THE INVENTION

#### Plant defense

15 Most plants are susceptible to infection by pathogens such as nematodes and develop various undesirable disease symptoms upon infection which cause retarded growth, reduced yield and consequently economical loss to farmers. The plants respond to infection with several defense mechanisms including production of phytoalexins, deposition of lignin-like material, accumulation of cell wall hydroxyproline-rich glycoproteins, expression of pathogenesis related proteins (PR-proteins) and an increase  
20 in the activity of several lytic enzymes. Some of these responses can be induced not only directly by infection but also in some cases by exposure to exogenous chemicals such as ethylene. The full capacity of the defense mechanism of the plant is, however, normally delayed in relation to the onset of infection, and thus, the plant may be severely injured before its defense mechanism reaches its maximum capacity. Also, the defense  
25 mechanism of the plant may not in itself be sufficiently strong to effectively combat the infectious organism. This is in particular true for cultivated plants which have often been cultivated with the aim of increasing the yield, decreasing the climate susceptibility, decreasing the nutrient demand etc. Therefore, a normal and necessary procedure is to treat infected plants or plants susceptible to infection with a chemical either as a prophylactic treatment or shortly after infection. The use of a chemical treatment is  
30 neither desirable from an ecological nor from an economic point of view. Another procedure to combat the infectious organism is crop rotation. However, this is not able to fully overcome the problem. It would therefore be desirable to be able to enhance the

defense of the host plant itself by introducing new and/or improved genes by genetic engineering. The advantageous effect of the latter strategy would be the immediate inhibition of a phytopathogenic attack, leading to a retarded epidemic establishment of the infecting organisms in genetically engineered plant crops and thus an overall reduction in the effect of the infection.

One of the phytopathogenic organisms which are most wide spread and which are pathogenic to potato are the potato cyst nematodes (PCN) *Globodera pallida* and *G. rostochiensis*. These nematodes cause considerable losses to potato crop growing, up to 10% of the annual yield world wide. Because cysts are very persistent to chemical treatment and can survive for several years in the soil, the use of nematicides and crop rotation are only moderately effective. The present invention circumvents these drawbacks in the control of PCN.

#### **Durability of PCN resistance**

The durability of the resistance is determined by the extent of variation at (a)virulence loci which occur among the pathogen biotypes and the ability of the pathogen to generate novel specificities at (a)virulence loci. For PCN, the variation at (a)virulence loci is for the majority determined by the original founders which have been introduced into Europe. PCN are endemic in the Andes region of South-America where they coevolved with their Solanaceous hosts. They are thought to have been introduced into Europe relatively recently, after 1850, together with collections of potato species which were imported for breeding purposes. Only a limited part of the genetic variation present in their centre of origin has been introduced into Europe (Folkertsma 1997). From the moment of their introduction onwards, the genetic variation in virulence within and between European nematode populations has been determined predominantly by 1) the genetic structure of the primary founders, 2) random genetic drift and 3) gene flow. Mutation and selection can be excluded as a driving force for the observed variation; the species produce only one generation in a growing season, their multiplication rate is low, the time between generations is 2 to 4 years in normal crop rotation and the active spread of the nematode is limited to several centimeters in the soil. It seems therefore highly unlikely that PCN populations have acquired other virulence characteristics than those already present at the moment of their introduction into Europe. Strategies to obtain broad spectrum resistance against PCN are therefore based on combining a



minimal number of genes with complementary or partially overlapping resistance spectra (Bakker *et al.*, 1993).

### Plant disease resistance genes

5 The majority of plant resistance (R-) genes are located in chromosomal bins containing other disease or insect resistance factors (reviewed in Crute and Pink, 1996). These resistance genes are dominantly inherited, are often involved in resistance processes which are characterized by a hypersensitive response (HR) and are members of multigene families hypothesized to have evolved from common ancestral genes. Most R-  
10 loci are characterized by the presence of DNA sequences encoding putative gene products that contain (1) a nucleotide binding site (NBS) and (2) a leucine rich repeat structure (LRR). These structural motifs are known to occur in a large number of resistance gene products; nearly 30 resistance genes from various species have now been cloned and with the exception of two (*Hm1* and *mlo*; Johal and Briggs, 1992; Büschges  
15 *et al.* 1997), these genes are thought to be components of signal transduction pathways (Baker *et al.* 1997). On the basis of the structural similarity within the motifs of these genes, it is hypothesized that resistance genes are evolutionarily related components of a recognition system (Staskawicz *et al.* 1995). However, outside these structural motifs, the nucleotide sequences of disease resistance genes are unrelated and several subclasses can  
20 be distinguished (Leister *et al.* 1998). Genes associated with resistance to nematodes in potato are likely to constitute a separate subclass of R-genes. However, the basic architecture hereof has not yet been uncovered. The isolation, characterization and functional analysis of these nematode R-genes remains to be done.

Clustering of R-loci in potato has been reported. One of the large R-loci clusters  
25 is on the short arm of potato chromosome 5. This cluster comprises at least five R-loci: *R1* associated with resistance to *Phytophthora infestans* (Leonards-Schippers *et al.* 1992), *Nb* associated with HR type resistance to potato virus X (de Jong *et al.* 1997), *Rx2* associated with an extreme type of resistance to PVX, and *Gpa* and *Grp1* associated with resistance to the PCN (Kreike *et al.* 1994; Rouppe van der Voort *et al.* 1998). The  
30 recently identified PCN R-locus *Gpa5* is also located within the *Grp1* region (Rouppe van der Voort and Van der Vossen; unpublished data). Additionally, *Gpa6* has been mapped to a region on chromosome 9 on which the homologous region in tomato, *Sw5*, conferring resistance to tomato spotted wilt virus, resides (Rouppe van der Voort and

Van der Vossen; unpublished data).

### The *Gpa2* locus

The *Gpa2* locus in potato has been found to be associated with resistance to *G. pallida* populations D383 and D372 (Arntzen et al. 1994). The presence of a single locus in potato which acts specifically to this small cluster of populations indicates that a gene-for-gene relationship underlies this plant-pathogen interaction (Roupe van der Voort et al. 1997; Bakker et al. 1993). Although, the *Gpa2* locus has previously been mapped on the short arm of chromosome 12 of potato (Roupe van der Voort et al. 1997a), thusfar no sequence data or precise location were known. The gene was never isolated and no indication as to whether this single sequence would suffice to provide resistance or reduce susceptibility to nematode infection was available.

### SUMMARY OF THE INVENTION

The present invention relates to a nucleic acid sequence providing resistance to infection by a phytopathogenic nematode of the *Globodera* species when introduced into a host plant, said host plant prior to introduction being susceptible to infection to the phytopathogenic nematode, said introduction occurring in such a way that said nucleic acid sequence is expressed in the host plant. Furthermore the invention relates to sequences which are homologous to the aforementioned sequence and which, when present in a plant, are able to render said plant resistant to infection by *Globodera* species. More specifically, a sequence according to the invention is preferably that of SEQ ID NO.1 or a homologue thereof. The PCN resistance locus *Gpa2*, when present in a plant such as *Solanum* spp., is capable of conferring to the plant anti-phytopathogenic activity in the form of resistance to *Globodera* species which are known to invade and damage the roots of Solanaceae. The invention relates to the *Gpa2* resistance gene of which the DNA sequence is disclosed herein.

The invention also relates to a product encoded by a nucleic acid sequence according to the invention, said product providing nematode resistance activity. Furthermore, the invention relates to genetic constructs, vectors, host cells such as bacterial strains, yeast cells and plant cells comprising a nucleic acid sequence according to the invention. In another aspect, the present invention relates to a genetically transformed plant, preferably of the family Solanaceae, especially a genetically

transformed potato plant. Suitably, in a host cell according to the invention, the expression product of the nucleic acid sequence according to the invention, said expression product providing the anti-nematode activity, is produced in an increased amount as compared to the untransformed host cell so as to result in an increased resistance to *Globodera* species. A process for producing a genetically transformed or transfected nematode resistant plant is additionally provided as is a process for isolating or detecting nucleic acid sequences according to the invention, providing nematode resistance of the aforementioned type. A process for diagnosing whether a plant is resistant to *Globodera* species and a process for providing resistance to *Globodera* species to plant material are also disclosed in the present invention. The invention also encompasses a process for producing a polypeptide providing the resistance and a nematocide composition providing said resistance. Antibodies to the polypeptide are also envisaged as embodiments of the invention as is the application thereof in a diagnostic kit for assessing whether a plant is resistant to the aforementioned nematodes. A diagnostic kit according to the invention may also comprise probes and/or primers specific for detection of a nucleic acid sequence providing the resistance.

The present invention relates to oligonucleotides corresponding to a part of a sequence according to the invention or being complementary thereto, with which homologous resistance genes can be identified that confer resistance to *Globodera* species.

## DETAILED DESCRIPTION OF THE INVENTION

### Definitions

The following definitions are provided for terms used in the description and examples that follow.

- *Nucleic acid*: a double or single stranded DNA or RNA molecule.

- *Oligonucleotide*: a short single-stranded nucleic acid molecule.

- *Primer*: the term primer refers to an oligonucleotide which can prime the synthesis of nucleic acid.

- *Homologous sequence*: a sequence which has at least 70%, preferably 75%, more preferably 80%, most preferably 85% or even 90% sequence identity with the nucleic acid of the invention, whereby the length of the sequences to be compared for nucleic

acids is at least 100 nucleotides, preferably 200 nucleotides and more preferably 300 nucleotides and for polypeptides at least 50 amino acid residues, preferably 75 amino acid residues and more preferably 100 amino acid residues. Homology between the sequences may be as defined and determined by the TBLASTN computer programme for nucleic acids or the TBLASTP computer programme for polypeptides, of Altschul *et al.* (1990), which is in standard use in the art, or, and this may be preferred, the standard program BestFit, which is part of the Wisconsin Package, Version 8, September 1994, (Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA, Wisconsin 53711). Alternatively, a homologous sequence refers to a nucleic acid which can hybridize under stringent conditions to the nucleic acid of the invention. Nucleic acid hybridization is a method for detecting related sequences by hybridization of single-stranded nucleic acid probes with denatured complementary target DNA on supports such as nylon membrane or nitrocellulose filters. Nucleic acid molecules that have complementary base sequences will reform the double-stranded structure if mixed in solutions under the proper conditions, even if the target nucleic acid is immobilized on a support. Stringent conditions refer to hybridization conditions which allow a nucleic acid sequence of at least 50 nucleotides and preferably about 200 or more nucleotides to hybridize to a particular sequence at about 65°C in a solution comprising approximately 1 M salt, preferably 6 x SSC or any other solution having a comparable ionic strength, and washing at 65°C in a solution comprising about 0.1 M salt, or less, preferably 0.2 x SSC or any other solution having a comparable ionic strength. These conditions allow the detection of sequences having about 90% or more sequence identity. The person skilled in the art will be able to modify hybridization conditions in order to identify sequences varying in identity between 50% and 90% or more. Binding of the single-stranded nucleic acid probe to a corresponding target nucleic acid may be measured using any of a variety of techniques at the disposal of those skilled in the art.

- *Promoter*: the term "promoter" is intended to mean a short DNA sequence to which RNA polymerase and/or other transcription initiation factors bind prior to transcription of the DNA to which the promoter is functionally connected, allowing transcription to take place. The promoter is usually situated upstream (5') of the coding sequence. In its broader scope, the term "promoter" includes the RNA polymerase binding site as well as regulatory sequence elements located within several hundreds of base pairs, occasionally even further away, from the transcription start site. Such regulatory sequences are, e.g.,

sequences which are involved in the binding of protein factors which control the effectiveness of transcription initiation in response to physiological conditions. The promoter region should be functional in the host cell and preferably corresponds to the natural promoter region of the *Gpa2* resistance gene. However, any heterologous promoter region can be used as long as it is functional in the host cell where expression is desired. The heterologous promoter can be either constitutive or regulatable. A constitutive promoter such as the CaMV 35S promoter or T-DNA promoters, all well known to those skilled in the art, is a promoter which is subjected to substantially no regulation such as induction or repression, but which allows for a steady and substantially unchanged transcription of the DNA sequence to which it is functionally bound in all active cells of the organism provided that other requirements for the transcription to take place is fulfilled. A regulatable promoter is a promoter of which the function is regulated by one or more factors. These factors may either be such which by their presence ensure expression of the relevant DNA sequence or may, alternatively, be such which suppress the expression of the DNA sequence so that their absence causes the DNA sequence to be expressed. Thus, the promoter and optionally its associated regulatory sequence may be activated by the presence or absence of one or more factors to affect transcription of the DNA sequences of the genetic construct of the invention. Suitable promoter sequences and means for obtaining an increased transcription and expression are known to those skilled in the art.

- *Terminator*: the transcription terminator serves to terminate the transcription of the DNA into RNA and is preferably selected from the group consisting of plant transcription terminator sequences, bacterial transcription terminator sequences and plant virus terminator sequences known to those skilled in the art.

- *Nematode*: plant parasitic roundworms of the genus *Globodera*, i.e. *Globodera pallida* and *G. rostochiensis*.

- *Nematode resistance*: to understand the nature of the activity of the *Gpa2* locus in connection with nematode resistance, a brief description of the histopathology of *Solanum* spp. infected with *Globodera* species is hereby given. The infective second-stage larvae hatch and emerge from the cysts and then migrate to and enter roots of susceptible (nonresistant) and resistant potato plants. Before feeding and developing in the root tissue, the nematode induces the formation of multinucleated syncytium. In susceptible potato plants, cessation of feeding by the mature nematode is followed by the

development of cysts breaking out of the root tissue but still clinging to the potato roots. The larvae may survive for a long period in the cysts. In the case of a nematode resistant plant, the number of cysts formed by the adult female nematodes is reduced whereby retardation of the growth of the potato plant is prevented. In accordance herewith, the

5 term "nematode resistance" denotes the characteristic activity in a plant ascribable to a resistance gene, i.e. the capability of the gene products to reduce or prevent the formation of cysts on the roots of plants in particular of Solanaceae like e.g. *Solanum* spp.

- *Gene*: the term "gene" is used to indicate a DNA sequence which is involved in producing a polypeptide chain and which includes regions preceding and following the  
10 coding region (5'-upstream and 3'-downstream sequences) as well as intervening sequences, the so-called introns, which are placed between individual coding segments (so-called exons) or in the 5'-upstream or 3'-downstream region. The 5'-upstream region comprises a regulatory sequence which controls the expression of the gene, typically a promoter. The 3'-downstream region comprises sequences which are involved in  
15 termination of transcription of the gene and optionally sequences responsible for polyadenylation of the transcript and the 3' untranslated region. The term "resistance gene" is a nucleic acid comprising a sequence as depicted in Fig. 3 (SEQ ID NO.3), or part thereof, or any homologous sequence.

- *Resistance gene product*: a polypeptide having an amino acid sequence as depicted in  
20 Fig. 3 (SEQ ID NO.1) or part thereof, or any homologous sequence exhibiting the characteristic of providing nematode resistance when incorporated and expressed in a plant.

### Scope of the invention

25 The present invention relates to a nucleic acid sequence providing resistance to infection by a phytopathogenic nematode of the genus *Globodera* when introduced into a host plant, said host plant prior to introduction being susceptible to infection with the phytopathogenic nematode, said introduction occurring in such a way that said nucleic acid sequence is expressed in the host plant. Furthermore the invention relates to  
30 resistance sequences which are homologous to the aforementioned sequence and which, when present in a plant, are able to confer to said plant resistance to infection by *Globodera* species. More specifically, a sequence according to the invention is suitably that of SEQ ID NO.1 or a homologue thereof. The PCN resistance locus *Gpa2*, when

present in a plant such as *Solanum* spp., is capable of conferring, to the plant, anti-phytopathogenic activity in the form of resistance to *Globodera* species which are known to invade and damage the roots of Solanaceae. The invention relates to the *Gpa2* resistance gene of which the DNA sequence is disclosed herein.

5 Homologues of the nucleic acid sequence of the abovementioned embodiment of the invention which also provide resistance to *Globodera* species, said homologues being nucleic acid sequences encoding the amino acid sequence of SEQ ID NO.1, are also within the scope of the invention. A homologue of the nucleic acid sequence according to the invention can suitably also provide the resistance when said homologue is a  
10 nucleic acid sequence exhibiting more than 70% homology at nucleic acid level with SEQ ID NO. 1. Alternatively the homologue is a nucleic acid sequence exhibiting more than 75% homology at nucleic acid level with SEQ ID NO. 1, preferably exhibiting more than 80% homology at nucleic acid level with SEQ ID NO. 1, more preferably exhibiting more than 85% homology at nucleic acid level with SEQ ID NO. 1. A  
15 homologue of the nucleic acid sequence according to the invention, said homologue providing the resistance, can also be a nucleic acid sequence exhibiting more than 90% homology at nucleic acid level with SEQ ID NO.1 and can even be a nucleic acid sequence exhibiting more than 95% homology at nucleic acid level with SEQ ID NO.1. A homologue also providing the resistance can be a nucleic acid sequence capable of  
20 hybridising under normal to stringent conditions to the nucleic acid sequence of SEQ ID NO. 1. Naturally another suitable embodiment of a homologue of the sequence according to the invention, also providing the resistance, can be a nucleic acid sequence encoding a deletion, insertion or substitution mutant of the amino acid sequence of SEQ ID NO.1. Such a homologue, also providing the resistance, can be a nucleic acid sequence  
25 encoding a deletion, insertion or substitution variant, preferably as occurs in nature, of the amino acid sequence of SEQ ID NO.1. A nucleic acid sequence according to the invention may in addition to any of the embodiments described above or any combinations thereof further comprise at least one intron. Suitable examples of introns and locations thereof are provided in SEQ ID NO.2. A suitable embodiment of the  
30 nucleic acid sequence according to the invention is the genomic insert of pBINRGH2 as disclosed in the examples. A nucleic acid sequence according to the invention is suitably identical to that present in the genetic material of a species of the Solanaceae family, preferably a species of the genus *Solanum*. More specifically, such sequences can be

found on and are preferably identical to those present in the genome of potato on chromosomes 4, 5, 7, 9, 11 or 12. More specifically, the nucleic acid sequence is identical to that present in the genome of potato at the *Gpa2* locus. Obviously, a fragment of any of the above mentioned embodiments exhibiting the characteristic of providing the resistance falls within the scope of the invention.

According to the present invention, a DNA region comprising the PCN R-locus *Gpa2* has been isolated from a potato plant harbouring a wild *Solanum* genomic introgression segment possessing resistance against nematode infection. This resistance, which appears to be very effective in PCN control, is not present in most cultivated potato cultivars. Therefore, one object of the present invention is to provide plants, specifically *Solanum* spp., which have the features of cultivated plants, with anti-phytopathogenic activity in the form of resistance to *Globodera* species. Thus the present invention relates to a DNA segment comprising the *Gpa2* locus of about 200 kb comprising one or several genes, the gene product or gene products thereof being capable of conferring to the plant resistance to nematodes of the *Globodera* species.

Another aspect of the present invention is a nucleic acid comprising the *Gpa2* resistance gene, the nucleic acid having the sequence of all or part of the sequence depicted in Fig. 3 (SEQ ID NO.3) or any homologous sequence, including (where appropriate) both coding and/or noncoding regions and providing nematode resistance upon expression thereof in a plant. In a preferred embodiment the *Gpa2* gene comprises the deduced coding sequence provided in Fig. 3 (SEQ ID NO.1) or any homologous sequence, preceded by a promoter region and followed by a terminator sequence.

As described in the invention, the nucleic acid sequence according to the invention possesses very valuable features with respect to anti-nematode activity. Thus, the DNA region comprising the nucleic acid sequence according to the invention encoding a polypeptide conferring/evoking the anti-nematode activity as defined above, can be used for the construction of genetically modified hosts having an increased resistance to nematodes as compared to untransformed hosts. The nucleic acid region according to the invention is thus capable of being inserted into the genome of a host plant, which in itself is susceptible to infection by a nematode, in such a way that the nucleic acid sequence is expressed, thereby conferring to the host plant resistance to infection by a phytopathogenic nematode. Thus, another aspect of the present invention relates to a genetic construct consisting of the nucleic acid sequence according to the



invention which genetic construct can then be used to genetically transform a host, e.g. a plant such as a cultivated plant, in such a way that it becomes resistant to nematodes.

A genetic construct comprising a nucleic acid sequence according to any of the embodiments described above, said sequence being operably linked to a regulatory region for expression, falls within the scope of the invention. Accordingly, the present invention relates to a genetic construct comprising

- 1) a promoter functionally connected to
- 2) a nucleic acid region as defined according to the present invention
- 3) a transcription terminator functionally connected to the nucleic acid sequence.

Preferably, the regulatory region of a genetic construct according to the invention is a *Gpa2* regulatory region. Such a regulatory region can by way of example correspond to that present in the sequence of nucleotides 1-4874 of SEQ ID NO.3. The regulatory region can suitably even correspond to that of nucleotides 1-4874 of SEQ ID NO.3. The regulatory region preferably comprises a promoter functionally connected to the nucleic acid sequence as defined in any of the embodiments above or in the examples, said promoter being able to control the transcription of said nucleic acid sequence in a host cell, preferably in a plant cell.

The genetic construct may be used in the construction of a genetically modified host in order to produce a host showing an increased anti-nematode activity and thus an increased resistance towards nematodes. It will be understood that a large number of different genetic constructs as defined above may be designed and prepared. Without being an exhaustive list, elements of the genetic constructs which may be varied are the number of copies of each of the nucleic acid sequences of the genetic construct, the specific nucleotide sequence of each of the nucleic acid sequences, the type of promoter and terminator connected to each nucleic acid sequence, and the type of any other associated sequences. Thus, genetic constructs of the present invention may vary within wide limits.

The invention also relates to DNA constructs comprising the regulatory sequences, and more preferably the promoter region of the *Gpa2* resistance gene in conjunction with a structural gene sequence heterologous to said regulatory sequences.

A vector which carries a nucleic acid according to any of the embodiments disclosed above or in the examples or a genetic construct according to any of the embodiments disclosed above or in the examples also falls within the scope of the

invention. Preferably such a vector is capable of replicating in a host organism. The vector may either be one which is capable of autonomous replication, such as a plasmid, or one which is replicated with the host chromosome such as a bacteriophage or integrated into a plant genome. For production purposes, the vector is an expression vector capable of expressing the nucleic acid sequence according to the invention in the organism chosen for the production. Suitable cloning vectors, transformation vectors, expression vectors, etc..., are well known to those skilled in the art. A vector according to the invention is constructed to function in a host organism selected from the group consisting of a micro-organism, plant cell, plant, seed, seedling, plant part and protoplast. A host cell capable of resulting in a plant is preferred and suitably the host organism is selected from the group consisting of a plant, plant cell, plant part, seed, seedling and protoplast.

In a still further aspect, the present invention relates to a host organism which carries and which is capable of replicating or expressing an inserted nucleic acid region of the invention. Such a host organism is preferably selected from the group consisting of a micro-organism, plant cell, plant, seed, seedling, plant part and protoplast, harbouring a vector and/or a genetic construct as defined above. The term "inserted" indicates that the nucleic acid region has been inserted into the organism or an ancestor thereof by means of genetic manipulation, in other words, the organism may be one which did not naturally or inherently contain such a nucleic acid region in its genome, or it may be one which naturally or inherently contains such a nucleic acid region, but in a lower number so that the organism with the inserted nucleic acid region has a higher number of such regions than its naturally occurring counterparts. The nucleic acid region carried by the organism may be part of the genome of the organism, or may be carried on a genetic construct or vector as defined above which is harboured in the organism. The nucleic acid region may be present in the genome or expression vector as defined above in frame with one or more second nucleic acid regions encoding a second gene product or part thereof so as to encode a fusion gene product. The organism may be a higher organism such as a plant, or a lower organism such as a micro-organism. A lower organism such as a bacterium, e.g. a gram-negative bacterium such as a bacterium of the genus *Escherichia*, e.g. *E. coli*, or a yeast such as of the genus *Saccharomyces*, is useful for producing a recombinant polypeptide as defined above. The recombinant production may be performed by use of conventional techniques, e.g. as described by Sambrook et

al. (1990). Also, the organism may be a cell line, e.g. a plant cell line. Most preferably, the organism is a plant, i.e. a genetically modified plant such as will be discussed in further detail below. As mentioned above, the genetic construct is preferably to be used in modifying a plant. Accordingly, the present invention also relates to a genetically transformed plant comprising in its genome a genetic construct as defined above. The genetically transformed plant has an increased anti-nematode activity compared to a plant which does not harbour a genetic construct of the invention, e.g. an untransformed or natural plant or a plant which has been genetically transformed, but not with a genetic construct of the invention. Normally a constitutive expression of the gene products encoded by the genetic construct is desirable, but in certain cases it may be preferable to have the expression of the gene products encoded by the genetic construct regulated by various factors, for example by factors such as temperature, pathogens, and biological factors. The genetically transformed plant is obtained by introducing the nucleic acid sequence according to the invention within the genome of said plant having a susceptible genotype to nematodes, using standard transformation techniques. It will be apparent from the above disclosure, that the genetically transformed plant according to the invention has an increased resistance to nematodes as compared to plants which have not been genetically transformed according to the invention or as compared to plants which do not harbour the genetic construct as defined above. In a further aspect, the present invention relates to seeds, seedlings or plant parts obtained by growing the genetically transformed plant as described above or by genetically transforming a plant cell and generating said part. It will be understood that any plant part or cell derivable from a genetically transformed host of the invention is to be considered within the scope of the present invention.

A process for producing a genetically transformed host organism having increased resistance to *Globodera* species as compared to the host organism prior to the transformation, said process comprising transferring a genetic construct and/or a vector according to any of the embodiments disclosed above and in the examples into the host organism so that its genetic material comprises the genetic construct and/or vector and subsequently regenerating the host organism into a genetically transformed plant part is also a part of the invention. The host organism may be selected from the group consisting of a plant cell, plant, seed, seedling, plant part and protoplast of the plant type to be rendered resistant and may subsequently be regenerated to a plant. Preferably, the

nematodes against which resistance is provided are selected from the group consisting of *Globodera* species, more specifically *Globodera rostochiensis* and *Globodera pallida*. The host organism which is to be transformed is selected from a plant type of the family Solanaceae, preferably a *Solanum* spp. Plants of the species *Solanum tuberosum*, comprising commercial potato cultivars, are preferred as this is a particular problem area for the commercial growers of such plants.

In accordance with well-known plant breeding techniques it will be understood that the production of a genetically transformed plant may be performed by a double transformation event (introducing the genetic construct in two transformation cycles) or may be associated with use of conventional breeding techniques. Thus, two genetically modified plants according to the present invention may be the subject of cross breeding in order to obtain a plant which contains the genetic construct of each of its parent plants.

Additionally, the present invention also relates to the resistance gene product which is encoded by the nucleic acid sequence according to the invention and which has the deduced amino acid sequence provided in Fig. 3 (SEQ ID NO.1). Thus a polypeptide having an amino acid sequence provided in SEQ ID NO.1 and also a homologue of said amino acid sequence, said homologue being a substitution, insertion or deletion mutant conferring nematode resistance against *Globodera* species, form embodiments of the invention. A polypeptide according to the invention is encoded by a sequence according to any of the embodiments described above or in the examples. A process for producing such polypeptides having an amino acid sequence provided in SEQ ID NO.1, or a homologue of said amino acid sequence, said homologue being a substitution, insertion or deletion mutant possessing resistance to *Globodera* species, said process comprising the expression of the nucleic acid sequence or genetic construct according to any of the embodiments according to the invention and optionally isolating said polypeptide, said expression occurring in a host organism according to the invention, is also covered by the invention. A process comprising an isolation step of the polypeptide in a manner known *per se* for polypeptide isolation from cell culture or from the host organism itself is also covered.

A nematicide composition comprising as active component a polypeptide according to the above or produced according to the process described or a host organism expressing such a polypeptide in a formulation suitable for application as

nematicide to a plant and optionally comprising other ingredients required for rendering the polypeptide suitable for application as a nematicide, also falls within the scope of the invention. Preferably such a nematicide composition comprises the polypeptide in a slow release dosage form. It is quite efficient if such a nematicide composition is formulated and packaged comprising instructions for application as nematicide.

Antibodies may be raised against any purified resistance gene product according to the invention by any method known to those skilled in the art (for an overview see "Immunology - 5th Edition" by Roitt, Male: Pub 1998-Mosby Press, London). Such antibodies can be used for the detection of the gene product.

Another aspect of the invention relates to nucleic acid sequences comprising at least 16 contiguous nucleotides corresponding to or complementary to the *Gpa2* sequence, with the proviso that when such a nucleic acid comprises part or all of the NBS encoding sequence, the nucleic acid also comprises at least one codon of a 5' and/or 3' overhanging portion corresponding to the respective 5' and/or 3' adjacent amino acids of parts of the NBS sequence of the *Gpa2*, with the following sequence, GGIGKTT or GGLPLA (see Table 4). Preferably, the *Gpa2* sequence is comprised within the sequence of SEQ ID NO.1, 2 or 3. The sequence length is preferably at least 50 nucleotides, preferably more than 100 nucleotides rendering it suitable for use as a probe in a nucleic acid hybridization assay. Oligonucleotides complementary to one strand of the *Gpa2* sequence or part thereof, can be used as labeled hybridization probes in a Southern hybridization procedure or as primers in an amplification reaction such as the polymerase chain reaction (PCR), for the screening of genomic DNA or cDNA, or constructed libraries thereof, for the identification and isolation of homologous genes. Homologous genes that are identified in this way and which encode a gene product that is involved in conferring reduced susceptibility or resistance to a plant against pathogens, such as nematodes of the genus *Globodera*, are comprised within the scope of the invention. Suitable embodiments can be selected from any of the sequences SEQ. ID. No.4, 5, 6 and/or 7.

A diagnostic kit for assessing the presence of nematode resistance in a plant to infection by a phytopathogenic nematode of the genus *Globodera*, said kit comprising at least one nucleic acid defined above as a probe or primer, for screening of nucleic acid from a plant or plant part to be tested and/or comprising an antibody as defined above, is also comprised within the scope of the invention.

The invention also covers a process for isolating or detecting a nucleic acid sequence according to the invention providing nematode resistance as described above and in the examples, said process comprising the screening of genomic nucleic acid of a plant with said nucleic acids or a fragment thereof as probe or primer, said probe or primer being at least 16 nucleotides in length, the identification of positive clones which hybridize to said probe or primer and the isolation of said positive clones and the isolation of the nucleic acid sequence therefrom. Such a process comprises screening genomic nucleic acid of a plant, preferably such a process comprises the screening of a genomic library of a plant with a nucleic acid sequence according to SEQ ID NO 3 or a fragment thereof as probe or primer, said probe being at least 16 nucleotides in length. Alternatively such a process comprises the screening of a cDNA library of a plant with the coding portion of a nucleic acid sequence according to the invention providing the nematode resistance, or a fragment thereof as probe or primer, said probe or primer being at least 16 nucleotides in length. Preferably, for the screening of a cDNA library of a plant, the coding portion of a nucleic acid according to SEQ ID NO.1 or a fragment thereof is used as probe or primer. The probe or primer can be comprised within the sequence of SEQ ID NO 1, SEQ ID NO 2 or SEQ ID NO 3. The above processes can use a nucleic acid amplification reaction such as PCR in conjunction with at least one primer corresponding to or being complementary to the nucleic acid sequence according to the invention providing the nematode resistance, or a fragment thereof, said primer being at least 16 nucleotides in length. The primer can be complementary to the nucleic acid sequence of SEQ ID NO.1, SEQ ID NO.2 or SEQ ID NO.3 or a fragment thereof, said primer being at least 16 nucleotides in length. A probe or primer in such a process comprises a nucleic acid sequence encoding the amino acid sequence of a part or all of the NBS sequence of *Gpa2*. Suitable examples of primers comprising a nucleic acid sequence encoding the amino acid sequence of a specific part or all of the NBS sequence of *Gpa2* are given below (see Table 4). For reasons of specificity, the process can comprise application of a primer comprising at least part of the NBS sequence of *Gpa2* and at least one codon of a 5' and/or 3' overhanging portion corresponding to the respective 5' and/or 3' adjacent amino acids of the previously specified NBS sequence of *Gpa2*. An example of such a primer comprises the specified part of the NBS sequence of *Gpa2* and at least one codon of a 5' and/or 3' overhanging portion corresponding to the respective 5' and/or 3' adjacent amino acids of the NBS sequence of *Gpa2* of SEQ ID

NO.1. Preferably, said primers correspond to a sequence selected from SEQ ID NO.4, SEQ ID NO.5, SEQ ID NO.6 and/or SEQ ID NO.7.

A process for diagnosing whether a plant is resistant to a phytopathogenic *Globodera* species, said process comprising the detection of the presence of a nucleic acid sequence providing nematode resistance as defined in any of the embodiments according to the invention, the presence of a genetic construct according to any of the embodiments according to the invention, the presence of a vector according to any of the embodiments according to the invention or the presence of a polypeptide as defined according to the invention, in the genetic material of plant material of a plant to be tested falls within the scope of the invention. Combinations of detection of the various elements are also covered. The nucleic acid sequence and the polypeptide being detected can be naturally present in the plant to be tested or can have been introduced via genetic engineering. A process for diagnosis according to the invention can comprise any of the nucleic acid sequence detection processes already described above as embodiments of the invention. More specifically the process can comprise applications of the diagnostic kit described according to the invention in an analogous manner to application of other nucleic acid assay kits comprising probes or primers or antibody known in the art. Suitably such a kit according to the invention will be provided with the appropriate instructions for application thereof. Amplification reactions of nucleic acid , use of probes in Southern analysis and use of antibodies in immunoassays are suitable examples of applications known in the art.

Another process within the scope of the invention is a process for providing resistance to a phytopathogenic *Globodera* species to plant material, said process comprising the introduction into the plant genome of a nucleic acid sequence providing nematode resistance as defined in any of the embodiments according to the invention, a genetic construct according to any of the embodiments according to the invention, a vector according to any of the embodiments according to the invention in the genetic material of plant material of a plant to be tested and thereby producing a transformed plant cell, plant propagating material, plant part or plant. Such introduction of genetic material should result in a transformed host with the introduced genetic material stably present in the host such that replication of said host is possible with said genetic material. Such a process may further comprise regenerating the resulting transformed or transfected plant cell, plant propagating material or plant part. The process of

introduction of the genetic material can occur as commonly described in the art for introduction of genetic material into the appropriate host type.

The nucleic acid sequence comprising the resistance as provided by the present invention has numerous applications of which some are described herein but which are not limiting to the scope of the invention.

The present invention is further described in detail below, whereby the map-based cloning strategy used to isolate the *Gpa2* resistance gene of the invention is explained. The strategy to isolate the *Gpa2* gene was as follows:

- 1) genetic fine mapping of the *Gpa2* locus;
- 2) construction of a BAC contig spanning the *Gpa2* locus;
- 3) identification of candidate resistance gene homologues (RGH);
- 4) complementation analysis.

The *Gpa2* locus was initially mapped on chromosome 12 using information on the genomic positions of 733 known AFLP markers (Roupe van der Voort *et al.*, 1997a and 1997b). By use of RFLP probes, *Gpa2* was mapped more precisely between markers GP34 and CT79 on the distal end of chromosome 12 (Roupe van der Voort *et al.*, 1997a), a 6 cM genetic interval that was previously shown to harbour the potato virus X (PVX) resistance gene *Rx1* (Fig. 1; Bendahmane *et al.*, 1997). Cosegregation of *Gpa2* and *Rx1* in the tetraploid *Rx1* mapping population (S1-Cara) and a diploid *Gpa2* mapping population (F1SHxRH) confirmed the assumed linkage between the two genes. The S1-Cara recombinants initially chosen to confirm this linkage delimited the *Gpa2* interval between markers IPM3 and IPM5 (Fig. 2; Bendahmane *et al.* 1997).

Fine mapping of the *Gpa2* locus was subsequently carried out using cleaved amplified polymorphic sequence (CAPS; Konieczny and Ausubel, 1993) markers derived from the IPM3-IPM5 interval, all of which were initially developed for the cloning of *Rx1* (Fig. 1). 2,788 S1-Cara genotypes were assayed for recombination events in the IPM3-IPM5 region. In addition, 598 F1SHxRH genotypes were subjected to a GP34/IPM5 marker screening as marker IPM3 was not informative in population F1SHxRH. Plants with recombination events between these markers were subsequently tested for all markers available in the IPM3-IPM5 region as well as for *Gpa2* resistance. This analysis showed that *Gpa2* is located between markers IPM4c and 111R (Fig. 2). Among the 2,788 S1-Cara genotypes and 598 F1SHxRH genotypes tested, only one genotype, S1-761, was identified in which a recombination event had occurred between



*Gpa2* and marker IPM4c (Fig. 2B). On the other side of *Gpa2*, genotype S1-B811 could be used to identify marker 111R as a flanking marker for the *Gpa2* interval (Fig. 2B).

Four BAC clones, BAC77, BAC45, BAC221 and BAC111, which map to the 0.06 cM IPM4c-111R genetic interval harbouring the *Gpa2* locus, were isolated from a BAC library prepared from a progeny of a selfed cv. Cara (Fig. 1C). However these four BAC clones did not completely cover the *Gpa2* interval. Screening of the Cara BAC library with CAPS markers 77R and 45L (Fig. 1B) did not lead to the identification of Cara BAC clones that spanned the region between markers 77R and 45L. A second BAC library was constructed from the diploid potato genotype SH83-92-488 (SH83). Screening of the SH83 potato BAC library with CAPS markers 77R and 45L did result in the identification of such a BAC clone (SHBAC43). In this way a contiguous physical map of the IPM4c-111R *Gpa2* interval was constructed comprising SHBAC43, BAC45, BAC221a and BAC111 (see Fig. 2C). Restriction analysis of this BAC contig delimited the physical size of the *Gpa2* locus of approximately 200 kb.

As the size of the *Gpa2* locus was still too large for direct localization of the *Gpa2* resistance gene by complementation analysis, BAC clones SHBAC43, BAC45, BAC221a and BAC111 were analysed for the presence of R-gene homologous sequences. Despite the general lack in DNA sequence conservation between R-genes, there are a few conserved protein motifs in the NBS region present in many of these genes. Leister *et al* (1996) has shown that it is possible to amplify resistance gene like sequences from potato using degenerate primers based on these homologous regions. Using degenerate primers RG1 and RG2 (Aarts *et al.*, 1998), whose sequences are based on the conserved P-loop and domain 5 region of the NBS in the N, L6 and RPS2 R-genes (Whitham *et al.*, 1994; Lawrence *et al.*, 1995; Bent *et al.*, 1994 and Mindrinis *et al.*, 1994) a DNA fragment of the expected size (approximately 530 bp) was amplified from BAC221a. Southern analysis of *Eco*RI restricted DNA of SHBAC43, BAC45, BAC221a and BAC111 using the amplified PCR fragment from BAC221a as a probe, identified two copies of this R-gene like sequence on SHBAC43, one single copy on BAC221a and one copy on BAC111 (Fig. 2D). Subsequent sequence analysis of the complete inserts of these BAC clones showed that the previously identified R-gene like sequences on the BAC clones belonged to putative resistance gene homologues (RGHs). Three of these RGH sequences were designated to be candidates for the *Gpa2* gene and selected for complementation analysis; RGH1 on BAC221a, RGH2 on BAC111 and

RGH3 on SHBAC43. A fourth RGH identified on SHBAC43 contained marker IPM4c and therefore lies outside of the *Gpa2* interval (see Fig. 2C and 2D).

Genomic fragments of approximately 11 kb, 10.3 kb and 5.5 kb harbouring RGH1, RGH2 and RGH3, respectively, were subcloned from the BAC inserts into the plant transformation vector pBINPLUS (Van Engelen *et al.*, 1995) and transferred to a susceptible potato genotype using standard transformation methods. Roots of *in vitro* grown primary transformants were tested for PCN resistance as described in Example 1. This *in vitro* resistance assay revealed that the 10.3 kb genomic insert harbouring RGH2 was able to complement the susceptible phenotype. RGH2 was therefore designated the *Gpa2* gene, the DNA sequence which is provided in Fig. 3.

The following examples provide a further illustration of the present invention which is nevertheless not limited to these examples.

## EXAMPLES

### EXAMPLE 1: ASSESSING NEMATODE RESISTANCE

#### A. *In vivo* resistance assay

Eggs and second stage juveniles ( $J_2$ ) are obtained by crushing cysts which have been soaked in tap water for one week. The egg/ $J_2$  suspension is poured through a 100  $\mu$ m sieve to remove debris and cystwalls. Before inoculation, three to four week old potato stem cuttings are transferred from a peat mixture to 900 gram pots containing a mixture of silversand and a sandy loam fertilized with Osmocote (N-P-K granulates). Subsequently, plants are inoculated with nematodes to a final density of 5 egg/ $J_2$  per gram soil. Of each plant genotype, three replicates per nematode source are inoculated. Six replicates of the parental clones as well as resistant and susceptible standards are included for resistance tests with each nematode source. Resistant standards are *Solanum tuberosum* cv. Multa (resistant to *G. pallida* D383), *S. vernei* hybrid 58.1642/4 (resistant to *G. rostochiensis* line Ro<sub>1</sub>-19) and *S. vernei* hybrid 62-33-3 (resistant to both D383 and Ro<sub>1</sub>-19). The susceptible standard is *S. tuberosum* cv. Eigenheimer. Plants are arranged in a randomized block design and grown in a greenhouse with 15°C and 25°C as minimum and maximum temperature, respectively.

After three months, cysts are recovered from the soil with a Fenwick can

(Fenwick 1940) and the size of the root systems is judged on a scale of 0 to 3. Resistance data of a genotype are only recorded when at least two well-rooted plants of this genotype are available. The mean cyst numbers developed per genotype are standardized using a  $\log_{10}(x + 1)$  transformation and then subjected to SAS Ward's minimum variance cluster analysis (SAS Institute Inc., Cary NC, USA). On the basis of this analysis the plants are divided into a resistant, an unassigned or a susceptible class.

#### B. *In vitro* resistance assay

Alternatively, the resistance assay is carried out on sterile tissue culture plants in agar. Two or three nodia from each *in vitro* grown (transgenic) potato plant are grown on solidified B5 medium (Gamborg *et al.* 1968) with 0.5% Phytigel™ (Sigma) and 2% sucrose for one week (25°C and 16 hr light regime). Each new root tip (on average 2 per nodium) is then inoculated with 15 sterilized second stage juveniles. Preparation of inoculum

is essentially as described by Heungens *et al.* (1995) with slight modifications. Cysts are collected in a modified 20 ml syringe with a 22 µm nylon mesh and surface sterilized in 90% ethanol for 15 sec followed by an 8 min wash in 1.3% (w/v) commercial bleach. To remove excess bleach, the cysts are washed three times in sterile tap water for 5 min and incubated in sterile tap water for 3 days at 20°C in the dark. Cysts are then transferred to filter sterile potato root differentiate (PRD) and left to hatch for 5 days at 20°C in the dark. Second stage juveniles are subsequently transferred to a 5 µm sieve-syringe and incubated first in 0.5% (w/v) streptomycine-penicilline G solution for 20 min, then in 0.1% (w/v) ampicillin-gentamycin solution for 20 min and finally in 0.1% chlorhexidin solution for 3 min. After three 5 min wash steps in sterile tap water the second stage juveniles are suspended in the required volume (sterile tap water) for inoculation. The petridishes with the inoculated root tips are incubated in the dark at 20°C. After 5-6 weeks the level of infection is determined by counting the number of female nematodes formed on the roots.

#### EXAMPLE 2: COSEGREGATION OF *Gpa2* NEMATODE RESISTANCE AND *Rx1* VIRUS RESISTANCE.

The *Gpa2* locus was initially mapped to chromosome 12 using information on the

genomic positions of 733 known AFLP markers (Roupe van der Voort *et al.*, 1997a and 1997b). By use of RFLP probes, *Gpa2* was mapped more precisely between markers GP34 and CT79 on the distal end of chromosome 12 (Fig. 2A; Roupe van der Voort *et al.*, 1997a), a 6 cM genetic interval that was previously shown to harbour the potato virus X (PVX) resistance gene *Rx1* (Bendahmane *et al.*, 1997).

To confirm the assumed linkage between *Gpa2* and *Rx1* (Roupe van der Voort *et al.* 1997), a pilot experiment was carried out in which the segregation of both genes was followed in two different mapping populations; a tetraploid ( $2n = 4x = 48$ ) mapping population derived from a selfing of potato cv. Cara (S1-Cara), initially constructed for fine mapping of *Rx1* (Bendahmane *et al.* 1997), and the diploid ( $2n = 2x = 24$ ) *Gpa2* mapping population derived from a cross between the diploid potato clones SH83-92-488 and RH89-039-16 (F1SHxRH; Roupe van der Voort *et al.*, 1997a and 1997b). Potato genotypes Cara and SH have the wild accession *Solanum tuberosum* spp. *andigena* CPC 1673 in common.

The tests for *Gpa2* and *Rx1* resistance were performed on the parental genotypes Cara, SH83 and RH89, four S1 genotypes which were recombined in a 1.21 cM interval between markers GP34 and IPM5 (Fig. 1B; Bendahmane *et al.* 1997) and two F1SHxRH genotypes which harboured cross-over events in a 6 cM interval between markers GP34 and CT79 (Roupe van der Voort *et al.* 1997). The PVX resistance assay was carried out using a cDNA of the PVX<sub>CP4</sub> isolate (Goulden *et al.* 1993). Potato plants were graft-inoculated with scions of *Lycopersicon esculentum* cvs. Ailsa Craig or Money Maker systemically infected with PVX<sub>CP4</sub>. Northern blots were prepared from total RNA isolated from newly formed potato shoots 3-4 weeks post-inoculation (Bendahmane *et al.* 1997). Extreme PVX resistance or susceptibility was determined by the presence or absence of a hybridization signal on Northern blots probed with <sup>32</sup>P-labelled cDNA of PVX<sub>CP4</sub> (Chapman *et al.* 1992). Three replicates per genotype were assayed. For the *Gpa2* assay *G. pallida* population D383 was used (Roupe van der Voort *et al.* 1997a). The nematode resistance assay was performed as described in Example 1A. Nematode population Rookmaker with different virulence characteristics as population D383 (Bakker *et al.* 1992) was used to confirm the specificity of *Gpa2* resistance in tested plants.

The resistance tests showed a clear reduction in the number of cysts of *G. pallida* population D383 on genotypes which were resistant to PVX<sub>CP4</sub>. The number of cysts

developed on the resistant S1-Cara genotypes appeared to be slightly higher than the number of cysts found on the resistant genotypes of population F1SHxRH. However, a considerable reduction in size of these cysts was observed as compared to the cysts developed on a susceptible genotype. This observation was corroborated after comparing the number of eggs per cyst developed on resistant and susceptible genotypes. Average cyst contents were determined from at least 30 cysts (if possible) and subjected to a *t*-test. A significant difference (at  $P < 0.05$ ) was found between the average number of eggs per cyst developed on Cara, SH83 and cv. Multa (resistant control), and average egg contents per cysts recovered from genotype S1-350, RH89 and cv. Eigenheimer (susceptible control). Resistance tests using *G. pallida* population Rookmaker show that cv. Cara is susceptible to this nematode population, indicating a specificity for the *G. pallida* resistance in population S1-Cara.

Although limited numbers of S1-Cara and F1SHxRH genotypes were tested for resistance to *G. pallida* population D383 and PVX respectively, based on the position of the crossover events in the tested plants it could be concluded that *Gpa2* and *Rx1* cosegregate in both mapping populations (with a maximum probability of  $P = 1/64$  that the observed linkage could be explained by chance). The tested S1-Cara recombinants were previously used to delimit the *Rx1* interval between markers IPM3 and IPM5 (Bendahmane *et al.* 1997). Cosegregation of *Gpa2* with *Rx1* indicates therefore that *Gpa2* also resides in this region (Fig. 2A).

**EXAMPLE 3: ISOLATION OF CARA BAC CLONES AND PRODUCTION OF CAPS MARKERS DERIVED FROM THE *Rx1/Gpa2* LOCUS** (according to the unpublished article in preparation of Kanyuka, K., Bendahmane, A., Rouppe van der Voort, J.N.A.M., van der Vossen, E.A.G. and Baulcombe, D.C. Mapping of intra-locus duplications and introgressed DNA: aids to map-based cloning of genes from complex genomes illustrated by analysis of the Rx locus in tetraploid potato).

#### *Construction of a Cara BAC library*

In order to clone the *Rx1* locus, a BAC library of 160,000 clones was prepared from plant SC-781 which is a progeny of selfed cv Cara carrying *Rx1* in the duplex condition (*Rx,Rx,rx,rx*). High molecular weight DNA was prepared in agarose plugs from potato protoplasts essentially as described in Bendahmane *et al.* (1997). The agarose plugs

were dialysed three times for 30 min against TE buffer (10 mM Tris-HCl pH 7.5, 1 mM EDTA), once at room temperature and twice at 4°C. The plugs were then equilibrated in *Hind*III buffer (10 mM Tris-HCl, 10 mM MgCl<sub>2</sub>, 50 mM NaCl, 1 mM DTT, pH 7.9) twice on ice for 1 h. Half of each plug (~5 µg of DNA) was transferred to a test tube containing 360 µl of *Hind*III buffer and 10-15 units of *Hind*III restriction enzyme. The enzyme was allowed to diffuse into a plug at 4°C for 1 h and the digestion was carried out at 37°C for 30 min. The reaction was stopped by adding 1 ml of 0.5 M EDTA and plugs were immediately loaded into a 1% low melting point agarose gel and subjected to contour-clamped homogeneous electric fields (CHEF; Chu, 1989) electrophoresis in a CHEF DR II system (Bio-Rad Laboratories, USA) in 0.5 X TBE buffer (45 mM Tris-borate pH 8.0, 1 mM EDTA) at 150 volts for 10 h at 4°C and constant pulse time of 5 sec or 8 sec. Compression zones containing the DNA fragments of 100 kb or 150 kb were excised from the gel and dialysed against 30 ml TE in a 15 cm Petri dish for 2 h at 4°C. Dialysed agarose slices were then transferred to a 1.5 ml test tube, melted at 70°C for 10 min and digested with 1 unit of GELASE (Epicentre Technologies, USA) per 100 mg of agarose gel for 1 h at 45°C.

The size selected potato DNA (25-50 ng) was ligated to 25-50 ng of *Hind*III-digested and dephosphorylated pBeloBAC11 (Shizuya *et al.*, 1992) using 400 to 800 units of T4 DNA LIGASE (New England BioLabs, USA) at 16°C for 24 hours in a total volume of 50 µl. The ligation products were dialysed against 1 X TE using 0.025 µm MF-MILLIPORE MEMBRANE FILTER (Millipore, UK) at 4°C for 2 h and 30 min at room temperature using the "drop dialysis" method of Maruzyk and Sergeant (1980).

Transformation of *E. coli* DH10B cells was carried out by electroporation using a BRL CEMI-PORATOR SYSTEM (Life Technologies Ltd, UK). To 20 µl of electro-competent cells, 0.5-3 µl of ligation mixture was added. After electroporation, *E. coli* cells were mixed with 1 ml SOC medium (0.5% yeast extract, 2% tryptone, 10 mM NaCl, 2.5 mM KCl, 10 mM MgCl<sub>2</sub>, 10 mM MgSO<sub>4</sub>, 20 mM glucose) and incubated at 37°C for 1 h with gentle shaking (80 rpm). The cells were spread on Luria broth (LB) agar plates containing chloramphenicol (12.5 lg/ml), 5-bromo-4-chloro-3-indolyl-β-D-galactoside (Xgal) (40 lg/ml) and isopropyl-1-thio-β-D-galactoside (IPTG) (0.12 mg/ml). Plates were incubated at 37°C for 24 hours. DNA from the compression zones of 100 and 150 kb led to clones with an average insert size of 100 kb and a transformation efficiency of approximately 1000 and 150 white colonies per 11 ligation mixture,

respectively. Approximately 92000 white colonies from these ligations were picked individually into 384 well microtiter plates (Genetix, UK) containing LB freezing buffer (36 mM  $K_2HPO_4$ , 13.2 mM  $KH_2PO_4$ , 1.7 mM citrate, 0.4 mM  $MgSO_4$ , 6.8 mM  $(NH_4)_2SO_4$ , 4.4 % V/V glycerol, 12.5  $\mu g/ml$  chloramphenicol in LB medium), grown overnight at 37°C and stored at -80°C. Another 100 bacterial pools containing ~500-1000 white colonies each (these pools also contained approximately 500-1500 blue bacterial colonies with an empty pBeloBAC11) were prepared by scraping the colonies from agar plates into the LB medium containing 18% glycerol and 12.5  $\mu g/ml$  chloramphenicol using a sterile glass spreader. These pools were also stored at -80°C.

***Screening of the Cara BAC library with markers IPM3, IPM4 and IPM5 and isolation of BAC clones derived from the Rx1/Gpa2 locus***

The Cara BAC library was initially screened with CAPS markers IPM3, IPM4 and IPM5 corresponding to the AFLP markers PM3, PM4 and PM5 flanking the Rx1 locus (Bendahmane *et al.*, 1997). This was carried out as follows. For the first part of the library of 92,160 clones stored in 384 well microtiter plates the plasmid DNA was isolated using the standard alkaline lysis protocol (Heilig *et al.*, 1997) from pooled bacteria of each plate to produce 240 plate pools. Aliquots of these plate pools were combined to prepare 26 'superpools' containing DNA from 9 plate pools, and one superpool containing DNA from 6 plate pools. To identify individual BAC clones carrying the CAPS markers the superpools and then the corresponding plate pools were screened. Once an individual plate had been identified the clones corresponding to each of the 24 columns of the positive plate were grown for 3-4 h at 37°C in LB medium and PCR was carried out on 3  $\mu l$  of bacteria. After identification of a positive column a colony PCR on each of the corresponding 16 colonies of this column was carried out leading to identification of a single positive BAC clone.

For the second part of the library, which is stored as one hundred pools of approximately 1000 clones, plasmid DNA was isolated from each pool of clones using the standard alkaline lysis protocol and PCR was carried out to identify positive pools. Bacteria corresponding to positive pools were diluted, plated on LB agar plates and subsequently colony hybridisation was carried out as described in Sambrook *et al.* (1989) using  $^{32}P$ -labelled DNA probes corresponding to the CAPS markers. PCR with the corresponding CAPS primers was used to distinguish between hybridising colonies

carrying the markers previously mapped to homologues located elsewhere in the genome and those derived from the *Rx1* locus.

Positive BAC clones were analysed by isolating plasmid DNA from 5 ml overnight cultures (LB medium supplemented with 12.5 mg/ml chloramphenicol) using the standard alkaline lysis miniprep protocol (Engebrecht *et al.*, 1997) and resuspended in 50 µl TE. Plasmid DNA (10 µl) was digested with *NotI* for 3 h at 37°C to free the genomic DNA from the pBeloBAC11 vector. The digested DNA was separated by CHEF electrophoresis in a 1% agarose gel in 0.5 X TBE at 4°C using a BIO-RAD CHEF DR II system (Bio-Rad Laboratories, USA) at 150 volts with a constant pulse time of 14 sec for 16 h.

Screening of the Cara BAC library with marker IPM3 identified three BAC clones: BAC167, BAC191 and BAC117, with potato DNA inserts ranging from 100 to 120 kb (Fig. 1C). *DdeI* digestion of the IPM3 DNA in these BAC clones and other potato DNA samples revealed that BAC117 carried the IPM3 allele that was linked in *cis* to *Rx1*. The other two BAC clones, BAC167 and BAC191, contained alleles from a corresponding region of the *rx* chromosomes. To identify the relative genome positions of these BAC clones, pairs of PCR primers were designed based on the sequence of the right and left ends of the insert. Inverse polymerase chain reaction (IPCR; Ochman *et al.*, 1990) was used to isolate the right and left end sequences of insert DNAs. BAC DNA was isolated and digested separately with *NlaIII*, *HpaII*, *MseI*, *HinP1I*, *PvuII*, *HaeIII* (for isolation of a left end sequence) or with *RsaI*, *SacI*, *EcoRI*, *HaeIII*, *MaeII*, *MseI*, *PvuII*, *HinP1I* (for isolation of a right end sequence) for 4 h at 37°C and recircularised by self ligation for 2 h at 20°C. Ligations were carried out using 5-50 ng of digested DNA and 5-10 units of T4 DNA LIGASE (Boehringer Mannheim, Germany) in a total volume of 100 µl. PCR amplification of the recircularised DNA was carried out using 3 µl of self-ligated DNA as the template. AB1 (5'-CCTAAATAGCTTGCGTAATCATG-3') and AB2 (5'-TGACACTATAGAATACTCAAGCTT-3') primers were used for PCR amplification of the left end sequence of insert DNA. AB3 (5'-CGACCTGCAGGCATGCAAGCTT-3') and AB4 (5'-ACTCTAGAGGATCCCCGGGTAC-3') primers were used for PCR amplification of the right end sequence of insert DNA. PCR conditions were as follows: 94°C for 15 sec, 60°C for 15 sec, 72°C for 90 sec - for 35 cycles. PCR products were digested simultaneously with *HindIII* and the restriction enzyme used in the preparation



of IPCR DNA template. The released insert ends were gel purified and cloned into pGEM-3Z(f+) (Promega, USA). Sequences of the clones containing ~1-2 kb inserts were determined using a 377 or 373 DNA SEQUENCING SYSTEM (Applied Biosystems, UK). PCR tests using the BAC DNAs as templates showed that the BAC clones identified with marker IPM3 overlapped in the order BAC167, BAC117, BAC191, *Rx1* (Fig. 1C). The 191L marker was separated from *Rx1* by only a single chromosomal recombination event (in plant S1-1146; Fig. 1B) in a mapping population of 1720 plants. In the same population, 117L and IPM3 markers were separated from *Rx1* by two and three recombination events respectively whereas the GP34 marker, present in BAC167, was separated from *Rx1* by thirteen recombinations (Fig. 1B). The BAC library did not contain additional BACs extending further towards *Rx1* from the 191L marker.

Screening of the Cara BAC library with IPM4, which mapped at 0.06 cM from *Rx1* on the side away from IPM3 (Bendahmane *et al.*, 1997), identified two clones: BAC73 and BAC111, with inserts of ~70 kb each (Fig. 1C). *TaqI* digestion of the IPM4 CAPS marker in these clones suggested that BAC111 was linked in *cis* to the *Rx1* locus but that BAC73 carries DNA insert from the *rx* chromosome. To determine the relative genome position of BAC111 and BAC73 PCR tests were performed using end sequence primers of these BAC clones (Table 1). These tests suggested that BAC73 overlaps with BAC111 and that 73L and 111L represent opposite ends of this set of overlapping BACs. Both 73L and 111L co-segregated with IPM4. In the initial mapping population of 1720 individuals, these markers were separated from *Rx1* by one recombination event (in individual S1-761; Fig. 1B) and it was not possible to determine directly which of these markers was physically closer to *Rx1*. Hence, to orientate these BACs relative to *Rx1*, the Cara BAC library was screened with CAPS markers 111L and 73L. The BAC library was also screened with the IPM5 CAPS marker which is on the same side of *Rx1* as IPM4, but further from *Rx1* (Bendahmane *et al.*, 1997). It was anticipated that BACs containing IPM5 would orientate the 111L and 73L markers relative to *Rx1*. These analyses identified BAC218, carrying an allele of IPM5 identified by *PstI* digestion, as being linked in *cis* to *Rx1* (Fig. 1C). The end sequences of BAC218 insert DNA were converted into the CAPS markers, 218L and 218R, and mapped genetically to the recombination events between GP34 and IPM5. Marker 218L was positioned 0.48 cM (recombination fraction: 8/1720) from *Rx1*, between IPM5 and CT129. The 218R marker was positioned between IPM4 and IPM5, 0.30 cM (recombination fraction: 5/1720) from

Rx1. A single BAC pool #29 was also identified which contains three markers: 218R, 73L and 111R. CAPS analysis revealed that each of these markers in the BAC pool #29 is represented by the allele linked in *cis* to Rx1. Hence, it was concluded that BAC pool #29 contains a single BAC clone, BAC29, with DNA insert linked in *cis* to Rx1.

5 Therefore, BAC29 provided a link between BAC218 and the IPM4 BAC contig and orientated the markers from the IPM4 contig in the following order: Rx1, 111L, IPM4, 73L (Fig. 1B).

By screening the BAC library with 111L allele-specific primers BAC221 was identified which carries an insert DNA of 40 kb and is linked in *cis* to Rx1. The left  
10 end sequence of BAC221 is located inside of BAC111 whereas the right end sequence of BAC221 extends further towards Rx1 (Fig. 1C). However the marker 221R co-segregated with IPM4 in the S1-Cara mapping population and was separated from Rx1 by the recombination event in plant S1-761 (Fig. 1B).

To extend the IPM4 contig further towards Rx1 the Cara BAC library was  
15 screened with 221R allele-specific primers which identified BAC45 which has an insert DNA of 40 kb and is linked in *cis* to Rx1. The right end sequence of BAC45 is located inside of BAC221, whereas the left end sequence of BAC45, 45L, extends further towards Rx1 (Fig. 1C). However, BAC45 does not contain Rx1 as the CAPS marker 45L is genetically separated from Rx1 by the recombination event in plant S1-761 (Fig. 1B).  
20 Additional PCR screening of the BAC library with the 45L marker failed to identify any new BAC clones therefore leaving a gap between the IPM3 and IPM4 BAC contigs (Fig. 1C).

Taking into account that disease resistance loci in plants are often highly complex with small families of resistance genes clustered within several dozen kilobases  
25 (Ellis *et al.*, 1995; Hulbert and Bennetzen, 1991; Jones *et al.*, 1994; Martin *et al.*, 1993; Witham *et al.*, 1994), a low stringency PCR screening assay was developed for the identification of duplicated sequences related to CAPS markers from the vicinity of Rx1 (IPM3-IPM5 interval). Pools of DNA from 20 resistant plants (R pool) and 20 susceptible plants (S pool) and the individual BAC clones from the IPM4 contig were  
30 used as templates for PCR amplifications. Primer annealing temperatures in PCR reactions was 5 to 10°C lower than in conditions originally developed for each CAPS primer pair (Table 1) so that amplification of related sequences, in addition to the original marker, could take place. The PCR products obtained with a number of tested

CAPS primer pairs were the same size as the products produced under high stringency conditions. However, digestion of these low stringency PCR products with either *TaqI*, *AluI* or *DdeI* restriction enzymes revealed several new DNA fragments that were not identified previously. These included fragments that were nonpolymorphic as well as fragments polymorphic between the R and S pools. Digestion of the low stringency IPM4 products from the R pool with *TaqI* identified the original IPM4 locus (designated IPM4a) in BAC111. There were also new IPM4 restriction fragments that had not been detected previously. One of these fragments (IPM4b) was nonpolymorphic in the R and S pools. This fragment originated from BAC221 as the *TaqI* restriction fragment of similar size was also detectable after digestion of the IPM4b allele derived from this BAC (Fig. 1B). A second new DNA fragment was polymorphic between R and S pools and was not detected after digestion of either IPM4a or IPM4b alleles derived from BAC111 and BAC221, respectively. This fragment cosegregated with *Rx1* in all the plants of the S1-Cara mapping population, including plant S1-761 and others with recombination events between GP34 and IPM5. This new IPM4 marker allele was designated IPM4c (see Fig. 1B).

Screening of the Cara BAC library with IPM4 primers using conditions for the detection of the IPM4c allele identified a new BAC clone, BAC77, with a DNA insert of approximately 50 kb (Fig. 1C). The end fragments of BAC77 DNA insert were cloned, sequenced and converted into the CAPS markers 77L and 77R. Marker 77L cosegregated with both IPM4-c and *Rx1* whereas 77R was separated from *Rx1* by one recombination event in the recombinant individual S1-761 (Fig. 1B; based on analysis of 1720 segregants).

**TABLE 1:** Primer sequences and thermal cycling conditions for CAPS markers in the *Gpa2-Rx* interval.

Marker	Primers	PCR conditions	Restriction enzyme
GP34	5'-CGTTGCTAGGTAAGCATGAAGAAG 5'-GTTATCGTTGATTTCTCGTTCCG	94°C 15s 62°C 15s 72°C 1 min 35 cycles	<i>TaqI</i>
IPM3	5'-AGTAGTTTCAGGCTAGTG 5'-CAACATCACTTGATCAGAC	94°C 15s 54°C 15s 72°C 1 min 35 cycles	<i>DdeI</i>
117L	5'-CCTAGCGTAGAGCGGTGTATCCA 5'-GTAGACATTTAATAATTCGTCG	94°C 15s 57°C 20s 72°C 2 min 35 cycles	<i>RsaI</i>
191L	5'-ACAAATTGTATAATTATAGTGATACG 5'-CAAGACATTAATTAACCAAACAATGG	94°C 15s 50°C 15s 72°C 2 min 35 cycles	<i>EcoRI</i>
77L	5'-GCTTCTAAACTCTAAATTCAGATTC 5'-CATGTGCGGACTCGTTCTTTGTAG	94°C 15s 64°C 15s 72°C 1 min 35 cycles	<i>AluI</i>

Marker	primers	PCR conditions	Restriction enzyme
IPM4	5'-GTACTGGAGAGCTAGTAGTGATCA 5'-GAACACCTTAATACTACACGCTGCAGG	94°C 15s 62°C 15s 72°C 2 min 35 cycles	<i>TaqI</i>
77R	5'-CTCGAGGGATTGAATCCAAATTAT 5'-GGAAGCAGAATACTCCTGACTACT	94°C 15s 66°C 15s 72°C 1 min 35 cycles	<i>HaeIII</i>
45L	5'-GGAGTCAATGCAGGGTCTATGGA 5'-CTCATTTGACACTTCTCGAACACA	94°C 15s 62°C 15s 72°C 1 min 35 cycles	allele specific
221R	5'-GCTTACATTTGCTCGAAGAAGCCAC 5'-CCTTAATAATCAATAGATTCAACTCG	94°C 15s 60°C 15s 72°C 1 min 35 cycles	allele specific
111R	5'-CCACTGTGTAAGGGTCAACTATAGTC 5'-GAGATGAAGATTTTCTTGTCTGATGG	94°C 15s 65°C 15s 72°C 1 min 30s 35 cycles	allele specific
73L	5'-CATTTCTGAATTGCTTCCGACTTC 5'-CCATGAAAATTGTTATCACTGAGGTC	94°C 15s 60°C 15s 72°C 1 min 35 cycles	<i>AluI</i>
218R	5'-GATTACAGTTGTGAATTAGTTCGGTA 5'-GCAACAGATATATTCCACTTACCATTC	94°C 15s 62°C 15s 72°C 1 min 30s 35 cycles	<i>AluI</i>

5

10

**EXAMPLE 4: FINE MAPPING OF THE *Gpa2* LOCUS**

Cosegregation of *Gpa2* and *Rx1* resistance in both the mapping populations initially used to map the two loci, F1SHxRH and S1-Cara, respectively, delimited the *Gpa2* locus to the IPM3-IPM5 interval (see Example 2). For fine-mapping of the *Gpa2* locus, a total of 2,788 S1-Cara genotypes were assayed for recombination events in the IPM3-IPM5 interval. In addition 598 F1SHxRH genotypes were subjected to a GP34/IPM5 marker screening as marker IPM3 is not informative in population F1SHxRH. The GP34 CAPS marker is derived from a sequenced insert of RFLP clone GP34. The CAPS marker screening provided a total of 20 recombinants in the S1-Cara population and 9 recombinants in the F1SHxRH population. These recombinants were subsequently tested for the presence of markers 77L, IPM4c, 77R, 45L, 221R, IPM4a, 111R, 73L and 218R, all of which are derived from the PM3-IPM5 interval (see Fig. 2B), as well as for *Gpa2* resistance. The *Gpa2* resistance test was carried out using *G. pallida* population D383 (Roupe van der Voort *et al.* 1997a). The nematode resistance assays were performed on plants derived from *in vitro* stocks, stem cuttings or tubers. *In vitro* plants were transferred from MS medium containing 3% saccharose to a mixture of silversand and sandy loam under a moist chamber for one week. Two to four weeks after planting, plants showing vigorous growth were inoculated with nematodes. Assays were further performed as described for stem cuttings and tubers as described in Example 1 and in Roupe van der Voort *et al.* (1997a). *G. pallida* Rookmaker with different virulence characteristics as *G. pallida* D383 (Bakker *et al.* 1992) was used to confirm the specificity of *Gpa2* resistance in tested plants.

This analysis showed that *Gpa2* is located between markers IPM4c and 111R (Fig. 2B). Among the 2,788 S1-Cara genotypes and 598 F1SHxRH genotypes tested, only one genotype, S1-761, was identified in which a recombination event had occurred between *Gpa2* and marker 77R. On the other side of *Gpa2*, genotype S1-B811 identified marker 111R as a flanking marker for the *Gpa2* interval.

Marker orders deduced from the analysis of F1SHxRH corresponded to those found in population S1-Cara. Estimates of recombination frequencies and their standard errors were calculated with the aid of the program Linkage-1 (Suiter *et al.* 1983) by choosing the appropriate genetic model for each cross. Data for the non-recombinant class of genotypes were set for either a 3:1 segregation ratio for population S1-Cara or a

1:1 segregation ratio for population F1SH×RH since only strongly skewed segregation ratios will influence estimates of recombination frequencies notably (Säll and Nilsson 1994; Manly 1994). A chi-square test was used to test for differences in recombination frequencies between the marker intervals. The chi-square test criterion was determined from the recombinant and non-recombinant classes for each marker interval. Differences (rejection of the null hypothesis) were significant when the test criterion was greater than the  $X^2_{[0.05]}$  value. Estimates of recombination frequencies deduced from both populations were merged to obtain an estimate of the average recombination value for each marker interval. The graphical genotypes (Young and Tanksley, 1992) shown in Fig. 1 display the boundaries of the *Gpa2* interval.

#### EXAMPLE 5: CONSTRUCTION OF A CONTIGUOUS BAC CONTIG SPANNING THE *Gpa2* LOCUS

Example 3 describes the preparation of a Cara BAC library from a progeny of a selfed cv. Cara and the identification and isolation of BAC clones BAC77, BAC45, BAC221 and BAC111, which map to the 0.06 cM IPM4c-111R genetic interval harbouring the *Gpa2* locus (Fig. 1C). Additional PCR screening of the Cara BAC library with markers 45L and 77R failed to identify any BAC clones that spanned the region between BAC77 and BAC45.

To bridge this gap between BAC77 and the IPM4 BAC contig (see Fig. 2C), a second BAC library was constructed from the diploid potato genotype SH83-92-488. High molecular weight potato DNA was prepared in agarose plugs from potato nuclei as described in Liu *et al.* (1994) with the following modifications. Plant nuclei were isolated by grinding leaf tissue (10 g) in liquid nitrogen, suspending the powder in 100 ml nuclei isolation buffer (10 mM Tris-HCl pH 9.5, 10 mM EDTA, 100 mM KCl, 0.5 M sucrose, 4 mM spermidine 1.0 mM spermine, 0.1% mercaptoethanol) and sequential filtering through one layer each of 280, 88, 55 and 20 µm nylon mesh. One-twentieth volume of isolation buffer supplemented with 10% Triton X-100 was added to the filtrate and left on ice for 15 min. The nuclei were pelleted at 4°C (in 50 ml screwcap tubes) at 2200 rpm for 10 min and resuspended with isolation buffer to a final volume of 1 ml. The nuclei were heated at 42°C for 1-2 min, mixed gently with an equal volume of 1.4% low-melting point inCert agarose (FMC) prepared in 10 mM Tris-HCl pH 9.5

and 10 mM EDTA and immediately poured into molds to form plugs (V=100 µl/plug). The agarose plugs were treated with lysis buffer (1% sarkosyl, 0.4 M EDTA pH 8.5, 0.2 mg/ml proteinase K and 3.8 mg/ml sodiumdisulfite) at 50 °C for 2 days with one change of lysis buffer. Proteinase K activity was inhibited by incubating the agarose plugs 12 hours at 50°C in T<sub>10</sub>E<sub>10</sub> buffer (10 mM Tris-HCl pH 8.0, 10 mM EDTA) supplemented with 40 µg/ml PMSF.

Restriction enzyme digestion of the agarose plugs and subsequent size selection was carried out essentially as described in Example 3, with the following modifications. Half of each plug (~10 µg DNA) was digested with 10 U of *Hind*III restriction enzyme for 1 h. Size selection was carried out in two steps. Partially digested *S. tuberosum* DNA was initially subjected to CHEF electrophoresis at 4°C in 0.5 X TBE using a linear increasing pulse time of 60-90 sec and a field strength of 6 V/cm for 18 hr. After electrophoresis, the lanes containing the lambda DNA ladder were removed and stained with ethidium bromide to locate the region of the gel containing potato DNA fragments ranging from 100 to 150 kb in size. This region was excised from the gel using a glass coverslip and subjected to a second size selection step in a 1% SeaPlaque (low-melting point) agarose gel (FMC). CHEF electrophoresis was carried out for 10 hr at 4°C using a field strength of 4 V/cm and a constant pulse time of 5 sec. The compression zone containing DNA fragments of 100 kb was excised from the gel as described above and dialysed against 50 ml TE for 2 hr at 4°C. Dialysed agarose slices were then transferred to a 1.5 ml Eppendorf tube, melted at 70°C for 5 min and digested with 1 unit of GELASE (Epicentre Technologies, USA) per 100 mg of agarose gel for 1 hr at 45°C.

Ligation of the size selected DNA to *Hind*III-digested and dephosphorylated pBeloBAC11 and subsequent transformation of ElectroMAX *E. coli* DH10B competent cells (Life Technologies, UK) with the ligated DNA was carried as described in Example 3, using the BioRad Gene Pulser for electroporation. Settings on the BioRad Gene Pulser were as recommended for *E. coli* by the manufacturer. Approximately 60,000 white colonies were picked individually into 384 well microtiter plates containing LB freezing buffer, grown at 37°C for 24 hr and stored at -80°C.

By screening the SH BAC library, as described in Example 3, with CAPS markers 77R and 45L BAC clone SHBAC43 was identified (see Fig. 2C). For further analysis of SHBAC43, plasmid DNA was isolated from 5 ml overnight cultures (LB



medium supplemented with 12.5 mg/ml chloramphenicol) using the standard alkaline lysis miniprep protocol (Engbrecht *et al.*, 1997) and resuspended in 50 µl TE. Plasmid DNA (10 µl) was digested with *NotI* for 3 h at 37°C to release the insert DNA from the pBeloBAC11 vector, and subsequently analysed by CHEF electrophoresis. Comparison of the electrophoretic mobility of the SHBAC43 insert with that of the lambda concatemer ladder (BioRad) lead to the conclusion that SHBAC43 contains a BAC insert of approximately 110 kb.

## EXAMPLE 6: IDENTIFICATION OF CANDIDATE RESISTANCE GENE HOMOLOGUES (RGH) WITHIN THE *Gpa2* LOCUS

### *Identification of candidate RGHs*

Screening of BAC clones SHBAC43, BAC45, BAC221a and BAC111 with degenerate primers RG1 and RG2 based on conserved motifs within the NBS of the cloned resistance genes RPS2, N and L6 (see ; Aarts *et al.*, 1998) resulted in the weak amplification of a 530 bp fragment from BAC221a. The use of this fragment as a probe to screen a Southern blot containing *EcoRI* digested DNA of SHBAC43, BAC45, BAC221a and BAC111 showed that SHBAC43 contained 2 copies of this sequence and that BAC clones BAC221a and BAC111 each contained one copy of this sequence.

### *Sequence analysis*

The DNA sequence of BAC clones SHBAC43, BAC221a and BAC111 was determined by shotgun sequence analysis. For each BAC clone a set of random subclones with an average insert size of 2 kb was generated. 10 µg of CsCl purified DNA was sheared for 6 seconds on ice at 6 amplitude microns in 200 µl TE using a MSE soniprep 150 sonicator. After ethanol precipitation and resuspension in 20 µl TE the ends of the DNA fragments were repaired by T4 DNA polymerase digestion at 11°C for 25 minutes in a 50 µl reaction mixture comprising 1x T4 DNA polymerase buffer (New England BioLabs, USA), 1mM DTT, 100 µm of all 4 dNTP's and 25 U T4 DNA polymerase (New England Biolabs, USA), followed by incubation at 65°C for 15 minutes. The sheared DNA was subsequently separated by electrophoresis on 1% SeaPlaque LMP agarose gel (FMC). The fraction with a size of 1.5-2.5 kb was excised from the gel and dialysed against 50 ml TE for 2 hr at 4°C. Dialysed agarose slices were then transferred

to a 1.5 ml Eppendorf tube, melted at 70°C for 5 min, digested with 1 unit of GELASE (Epicentre Technologies, USA) per 100 mg of agarose gel for 1 hr at 45°C, and the DNA was subsequently precipitated. The 1.5-2.5 kb fragments were ligated at 16°C in a *EcoRV* restricted and dephosphorylated pBluescript SK<sup>+</sup> vector (Stratagene Inc.). The ligation mixture was subsequently used to transform ElectroMAX *E. coli* DH10B competent cells (Life Technologies, UK) by electroporation using the BioRad Gene Pulser. Settings on the BioRad Gene Pulser were as recommended for *E. coli* by the manufacturer. The cells were spread on Luria broth (LB) agar plates containing ampicillin (100 µg/ml), 5-bromo-4-chloro-3-indolyl-β-D-galactoside (Xgal) (64 µg/ml) and isopropyl-1-thio-β-D-galactoside (IPTG) (32 µg/ml). Plates were incubated at 37°C for 24 hours. Individual white colonies were grown in 96-well flat-bottom blocks (1.5 ml Terrific Broth medium containing 100 µg/ml ampicillin).

Plasmid DNA was isolated using the QIAprep 96 Turbo Miniprep system in conjunction with the BioRobot<sup>TM</sup> 9600 (QIAGEN) according to the manufacturers instructions. The ABI PRISM dye terminator cycle sequencing ready kit was used to perform sequencing reactions in a PTC-200 Peltier Thermal Cycler (MJ Research). The DNA sequences of the clones were determined using standard M13 forward and reverse primers. Sequence assembly was done using the 1994 version of the STADEN sequence analysis programme (Dear and Staden, 1991).

Analysis of the contiguous sequence of each BAC clone using the computer programme GENSCAN (Burge and Karlin, 1997) and BLASTX (Altschul *et al.*, 1990) identified a total of four NBS/LRR containing genes, two on SHBAC43, one on BAC221a and one on BAC111 (Fig. 2D). Three of these sequences were designated candidates for the *Gpa2* gene and selected for complementation analysis; RGH1 on BAC221a, RGH2 on BAC111 and RGH3 on SHBAC43 (Fig. 2D). The second NBS/LRR gene identified on SHBAC43 contained marker IPM4c and is therefore outside of the *Gpa2* interval (Fig. 2D).

#### EXAMPLE 7: TRANSFORMATION

For complementation analysis a 5.5 kb *SstI-XbaI* genomic fragment containing RGH3 from SHBAC43 and two *XbaI-XbaI* genomic fragments of approximately 11 kb and 10.3 kb containing RGH1 or RGH2 from BAC221a and BAC111, respectively, were

subcloned into the plant transformation vector pBINPLUS (Van Engelen *et al.*, 1995). These binary plasmids, designated pBINRGH1-3 were transferred to *Agrobacterium tumefaciens* strain AGL0 (Lazo *et al.*, 1991) by electroporation using the BioRad Gene Pulser. Settings on the BioRad Gene Pulser were as recommended for *A. tumefaciens* by the manufacturer. The cells were spread on Luria broth (LB) agar plates containing kanamycin (100 mg/L) and rifampicin (50 mg/L). Plates were incubated at 28°C for 48 hours. Small-scale cultures from selected colonies were grown in LB medium containing kanamycin (100 mg/l) and rifampicin (50 mg/l). Plasmid DNA was isolated as described previously and the integrity of the plasmids was verified by restriction analysis upon reisolation from *A. tumefaciens* and subsequent transformation to *E. coli*. *A. tumefaciens* cultures harbouring a plasmid with the correct DNA pattern were used to transform a susceptible potato genotype.

Transformation of the susceptible potato genotype, clone V, was essentially performed as described by Visser (1991) and is described briefly below. Stem explants (1 cm long internodes) were prepared from 5 week old tissue culture plants and precultured for 24 hours (25°C, 16 hour light regime) in Petri dishes containing 2 sterile filter papers saturated with PACM (feeding layers: MS30 medium supplemented with 2 g/l caseinehydrolysate, 1 mg/l 2,4 D and 0.5 mg/l kinetine, pH 5.8) which were placed on R3B medium (MS30 medium supplemented with 2 mg/l NAA and 1 mg/l BAP, pH 5.8). The explants were then infected for 10 minutes with an overnight culture of *A. tumefaciens* strain AGL0 containing either pBINRGH1, pBINRGH2, pBINRGH3 or the pBINPLUS vector, blotted dry on sterile filter paper and cocultured for 48 hours on the original feeder layer plates. Culture conditions were as described above. Overnight *A. tumefaciens* cultures were pelleted and resuspended in liquid MS20 medium prior to infection. Following cocultivation, the explants were transferred to MS20 medium (pH 5.8) supplemented with 1 mg/l zeatin, 200 mg/l cefotaxime, 200 mg/l vancomycin and 100 mg/l kanamycin and cultured under the culture conditions described above. The explants were transferred to fresh medium every 3 weeks. Emerging shoots were isolated and transferred to glass jars with selective medium lacking zeatin for root induction. Only transgenic shoots were able to root on the kanamycin containing medium.

**EXAMPLE 8: COMPLEMENTATION ANALYSIS**

*In vitro* grown transgenic ( $R_0$ ) plants were initially subjected to the *in vitro* resistance assay as described in Example 1B whereby sterilized second stage juveniles of *G. pallida* population D383 were used as inoculum. Three nodes from four independent primary transformants of the 4 different transformations were assayed;  $R_0$ (RGH1),  $R_0$ (RGH2) and  $R_0$ (RGH3) transgenic plants contain the candidate genes RGH1, RGH2 and RGH3, respectively, and  $R_0$ (BINPLUS) transgenic plants are without insert DNA and function as control plants. In addition, three nodes from 12 *in vitro* grown resistant and 12 *in vitro* grown susceptible progeny plants derived from the F1SHxRH mapping population (see Example 2) were included in the assay. The results are shown in Table 2. The development of nematode females on the roots of  $R_0$ (RGH1),  $R_0$ (RGH3) and  $R_0$ (BINPLUS) plants was similar to that observed on the roots of the susceptible control plants. In contrast, the  $R_0$ (RGH2) plants showed the same incompatible interaction with *G. pallida* population D383 as the resistant control plants. Three lines of evidence indicate that the 10.3 kb DNA fragment, which is integrated in the genome of  $R_0$ (RGH2) plants, harbours the *Gpa2* gene. First, the number of females able to develop on the roots of  $R_0$ (RGH2) plants was equivalent to the number of females able to develop on the roots of resistant control plants. Second, 90% of all the females that developed on these plants remained small and were translucent. This stagnation of female development was also observed on the roots of the resistant control plants. And third, the change in sex ratio (male/female=0.9) which is characteristic for the *Gpa2* phenotype was also observed for the  $R_0$ (RGH2) plants.

**TABLE 2** . Results of the complementation assay for *Gpa2* resistance.

Genotype	Average no. cysts/3 plants (# genotypes ) <sup>1)</sup>	Cyst phenotype
Susceptible F1SHxRH ( <i>gpa2</i> )	42 (12)	White
Resistant F1SHxRH ( <i>Gpa2</i> )	5 (12)	Translucent
R <sub>0</sub> (BINPLUS)	33 (4)	White
R <sub>0</sub> (RGH1)	39 (4)	White
R <sub>0</sub> (RGH2)	2 (4)	Translucent
R <sub>0</sub> (RGH3)	40 (4)	White

<sup>1)</sup> The numbers between brackets indicate the numbers of genotypes tested

#### **Molecular and computer analysis of the genomic insert conferring resistance**

To confirm the presence of the RGH2 insert in the R<sub>0</sub>(RGH2) with the resistant phenotype, a marker analysis with CAPS marker IPM4 was performed. The presence of the RGH2 linked CAPS marker IPM4a in all the R<sub>0</sub>(RGH2) plants transformed with pBINRGH2 indicates that the RGH2 gene is present in all these transgenic plants. Correct integration of the genomic fragment was also confirmed by Southern analysis using RGH2 and NPTII specific probes.

The sequence of the 10.3 kb *Xba*I-*Xba*I insert of pBINRGH2 is provided in Fig. 3C (SEQ ID NO.3). The genomic structure of the *Gpa2* gene was determined by analysis of the genomic sequence, derived from the insert of pBINRGH2, and cDNA fragments generated by 5' and 3' rapid amplification of cDNA ends (RACE). Comparison of these sequences revealed that *Gpa2* contains two introns at the C-terminus of the gene (Fig. 3). The first intron (237 bp) is located within the coding region of the gene whereas the second intron (112 bp) is situated within the 3'-untranslated region (Fig. 3). The second exon of *Gpa2* encodes a TGA stop codon and contains only 25 bp of coding nucleotides.

The deduced open reading frame (ORF) of the *Gpa2* gene encodes a predicted polypeptide of 912 amino acids with a MW of 104.5 KDa (SEQ ID NO.1).

**EXAMPLE 9: IDENTIFICATION AND MAPPING OF HOMOLOGOUS GENES.**

Screening of the SH83 BAC library as described in Example 4 using primers described in Leister *et al.* (1996) based on conserved motifs within the nucleotide binding site (NBS) of the cloned resistance gene RPS2 (GGVGKTT in case of primer S1 and GGLPLAL in case of primer AS1; see Tables 3 and 4) resulted in the amplification of DNA fragments of the expected sizes from all 156 BAC pools. This indicates that sequences homologous to the resistance gene motifs used to design primers S1 and AS1 are abundantly present in the potato genome.

Based on the nucleotide sequence of the resistance gene homologues RGH1-4, primers were designed for specific amplification of nucleic acid sequences cognate to the NBS of RGH1-4 (primers RG3 and RG4; see Tables 3 and 4). The position of primer RG3 corresponds to nucleotides 514-533 of SEQ ID NO.1 (Fig. 3). Primer RG4 is complementary to nucleotides 985-1002 of SEQ ID NO.1 (Fig. 3). These primers differ from RG1 and RG2 and those designed by Leister *et al.* (1996) in that the 3' terminal nucleotides are designed on the basis of amino acid residues that exceed the conserved residues used for the design of the former primers (see Table 4). PCR using primers RG3 and RG4 on template DNA of the BAC clones SHBAC43, BAC45, BAC221a and BAC111 resulted in amplification products of the expected size from SHBAC43, BAC221a and BAC111.

Screening of the SH83 BAC library as described in Example 4 using primers RG3 and RG4 identified 19 individual BAC clones that showed amplification of DNA fragments of the expected size. This indicates that these primers discriminate between RGH1-4 homologues and sequences containing common resistance gene motifs.

Primer sequences RG5 and RG6 (see Table 3) were designed on the basis of sequences outside of the NBS of RGH1-4. The position of primer RG5 corresponds to nucleotides 199-221 of SEQ ID NO.2 (Fig. 3). Primer RG6 is complementary to nucleotides 2681-2701 of SEQ ID NO.2 (Fig. 3). Screening the SH83 BAC library as described in Example 4 resulted in the isolation of 5 BAC clones which already were identified with primers RG3 and RG4. These BAC clones showed overlap with clones SHBAC43, BAC221a and BAC111. The primers RG5 and RG6 therefore discriminate between RGH sequences derived from the *Gpa2* locus and homologous variants

elsewhere on the potato genome. Primers RG3, 4, 5, 6 are SEQ ID NO. 4, 5, 6 and 7 respectively.

Mapping of the *Gpa2* homologues identified with primers RG3 and RG4 is carried out by developing CAPS markers designed on the end sequences of each BAC insert. These CAPS markers are used to screen 136 genotypes of population F1SHxRH. The data on marker segregation are scored and the respective loci are mapped on the SH83 genome by use of the computer package JoinMap2.0 (Stam, 1993). It is likely that one or more of these homologues map to regions of the potato genome harbouring mono- or polygenic resistance loci that confer resistance to other *G. pallida* or *G. rostochiensis* populations; such as *H1* (Pineda *et al.* 1993; Gebhardt *et al.* 1993), *Gpa* (Kreike *et al.* 1994), *Gpa5* (Roupe van der Voort and van der Vossen; unpublished data) and *Grp1* (Roupe van der Voort *et al.* 1998) on chromosome 5; *Gro1* on chromosome 7 (Barone *et al.*, 1990; Ballvora *et al.*, 1995); *Gpa6* on chromosome 9 (Roupe van der Voort and van der Vossen; unpublished data) and *Gpa3* on chromosome 11 (P. Wolters, unpublished data).

**Table 3:** Primer sequences and thermal cycling conditions for identification of *Gpa2* homologues

Primer	Primer sequence <sup>1)</sup>	PCR conditions	Expected product size
s1 as1	5'-GGTGGGGTTGGGAAGACAACG 5'-TGCTAGAGGTAATCCTCC	94°C 30s 51°C 30s 72°C 2 min 35 cycles	500 bp
RG1 RG2	5'-GGIATGGGIGGIGTIGGIAARACNACN 5'-ICCIAGIACYTTIARIGCIARIGGIARWCC	94°C 30s 50°C 30s 72°C 2 min 30 cycles	530 bp
RG3 RG4	5'-GGAGGCATCGGGAAAACAAC 5'-TGCTAGAGGTAACCCTCC	94°C 30s 55°C 30s 72°C 2 min 30 cycles	488 bp
RG5 RG6	5'-GATATGGTTGACTCGGAATCAAG 5'-GAGTATGGACCTCGATAGAGC	94°C 30s 60°C 30s 72°C 3 min 30 cycles	2500 bp

<sup>1)</sup> R=A or G; Y=T or C; W=A or T



**TABLE 4.** Oligonucleotides based on conserved peptide motifs within the NBS of PPS2 and RGHS

Motif / primer	Primer designation	Sequence <sup>2)</sup>
P-loop (RPS2/N/L6) s1	sense	G G V G K T T ggt ggg gtt ggg aag aca acg
P-loop (RGH1-4) RG3	sense	G G <u>I</u> G K T T gga ggc <u>atc</u> ggg aaa aca ac
GLPLAL (RPS2/N/L6) as1	antisense <sup>1)</sup>	G L P L A L caa cgc tag tgg caa tcc
GGLPLA (RGH1-4) RG4	antisense <sup>1)</sup>	G G L P L A <u>tgc</u> tag <u>agg</u> <u>taa</u> <u>ccc</u> <u>tcc</u>

<sup>1)</sup> Antisense primers are written in opposite orientation to the peptide sequence

<sup>2)</sup> Differences between primers s1/as1 and primers RG3/RG4 are underlined

#### EXAMPLE 10: A MARKER ASSISTED SELECTION ASSAY FOR *Gpa2*

The *Gpa2* locus is hypothesized to be introgressed from *S. tuberosum* spp. *andigena* CPC1673 into European cultivars. Flanking markers tightly linked to *Gpa2* are likely to be diagnostic for the presence of *Gpa2* in these cultivars. Therefore, *Gpa2* linked CAPS markers were used to screen two clones (abbreviated as CPC1673-a and CPC1673-b) of the wild species *Solanum tuberosum* spp. *andigena* CPC 1673 (hereafter referred to as CPC1673) as well as nine potato cultivars harbouring introgressions from CPC1673. The CAPS marker profiles were highly similar for the selfed CPC1673 genotypes and the analyzed potato cultivars harboring introgressions from CPC1673. The CAPS marker alleles linked to *Gpa2* were only identified in regions which appeared to be of CPC1673 origin. Among the seven CPC1673 cultivars tested, five differences in the size of an

introgressed region of 0.9 cM were observed. All *Gpa2* containing cultivars harbored the *Gpa2* flanking markers 77R and 111R thereby demonstrating that these markers are indicative for the presence *Gpa2* (see Table 5).

**TABLE 5:** Potato clones having *S. tuberosum* spp. *andigena* CPC1673 in their pedigree (with the exception of clone RH89) tested on the presence of chromosome 12 specific CAPS alleles. Resistance or susceptibility to *G. pallida* population Pa2-D383 is indicated by "R" or "S" respectively. Presence or absence of a CAPS marker band that cosegregates with resistance in populations S1-Cara and F1SHxRH is indicated by either a "+" or a "-". The order of the presented CAPS markers corresponds to the marker order on chromosome 12.

Clone	<i>Gpa2</i>	IPM3	191L	77L	IPM4c	77R	IPM4	111R	73L	218R	IPM5
CPC1673-a	n.d.	+	+	+	+	+	+	+	+	+	+
CPC1673-b	n.d.	+	+	+	+	+	+	+	+	+	+
Cara	R <sup>a)</sup>	+	+	+	+	+	+	+	+	+	+
Alcmaria	R <sup>b)</sup>	-	+	+	+	+	+	+	+	+	+
Multa	R <sup>a)</sup>	-	-	+	+	+	+	+	+	+	+
SH83	R <sup>a)</sup>	-	-	-	+	+	+	+	+	+	+
Amaryl	R <sup>b)</sup>	-	-	-	+	+	+	+	+	+	+
Marijke	R <sup>b)</sup>	-	-	-	+	+	+	+	+	+	+
Saturna	S <sup>a)</sup>	-	-	-	-	-	-	-	-	-	+
RH89	S <sup>a)</sup>	-	-	-	-	-	-	-	-	-	-

<sup>a)</sup> As determined by cyst counts on at least three replicates

<sup>b)</sup> Data from Arntzen et al. (1994)

## FIGURES

**Fig. 1.** High resolution map of the *Rx* locus (not drawn to scale). **A.** Simplified genetic map of potato chromosome XII (denoted by a horizontal line) in which the area left of the arrow is reversed between the potato and tomato genetic maps (Tanksley *et al.*, 1992). Vertical lines indicate positions of previously mapped RFLP markers (Bendahmane *et al.*, 1997; Tanksley *et al.*, 1992). The filled rectangle denotes a genetic interval between markers GP34 and 218L, which is magnified in panels B and C. **B.** Genetic map of the GP34-218L interval (denoted by a horizontal line). Positions of the RFLP marker GP34 and the AFLP markers IPM3, IPM4a and IPM5 are indicated by vertical lines. The positions of BAC end-derived markers and low-stringency PCR markers (enclosed in square brackets) are indicated by vertical arrows. The symbols L and R denote the BAC ends that were mapped relative to *Rx1*. The numbers in brackets below the bar indicate the numbers of S1-Cara individuals containing recombination events in each marker interval, identified in the initial S1-Cara mapping population of 1720 individuals. The predicted position of *Rx1*, delimited by the cross-over events in plants S1-1146 and S1-761, is indicated by the horizontal arrow. **C.** Positions of Cara BAC clones in the GP34-218L interval. Each open rectangle represents one BAC insert DNA. Inside of each rectangle is the name of the BAC clone, the estimated insert size in kb (except for the BAC29).

**Fig. 2.** High resolution genetic and physical map of the *Gpa2* locus. **A.** Relative position of the *Gpa2* locus on chromosome 12 of potato. Vertical lines indicate positions of previously mapped RFLP markers. The filled rectangle denotes the *Gpa2* locus between markers IPM3 and IPM5 which is magnified in panel B. **B.** High resolution genetic map and graphical genotypes of the IPM3-IPM5 interval, indicating differences in the size of *Solanum tuberosum* spp. *andigena* CPC1673 derived segments in different potato genotypes. The relative positions of CAPS markers used to fine-map *Gpa2* are indicated by vertical bars. The presented genotypes border the *Gpa2* interval. Introgression segments are indicated by thick bars. Size of marker intervals are not drawn to scale. The symbols R (for resistant) and S (for susceptible) indicate the *Gpa2* phenotype of the tested genotypes. **C.** High resolution physical map

of the *Gpa2* locus. The relative positions of CAPS markers are indicated by vertical bars. The open rectangles represent BAC clones isolated from the Cara BAC library. The shaded rectangle represents a BAC clone isolated from the SH83 BAC library. The name of each BAC clone is depicted within the rectangle and the estimated insert size is given in kb. The predicted position of *Gpa2* is indicated by the horizontal arrow. Recombinant S1-Cara genotypes S1-761 and S1-B811 delimit the *Gpa2* genetic interval. **D.** Relative positions of four resistance gene homologues (RGH1-4) identified within the IPM4c-111R *Gpa2* interval.

**Fig. 3.** Nucleic and amino acid sequence of the *Gpa2* gene. **A.** Coding nucleic acid sequence and deduced amino acid sequence of the *Gpa2* gene. **B.** Coding sequence of the *Gpa2* gene including intron 1. The position of intron 1 is indicated in bold italics (position 2712-2948). **C.** Sequence of the 10.3 kb *XbaI-XbaI* genomic DNA fragment inserted in pBINRGH2, harbouring the *Gpa2* gene. The initiation ATG codon (position 4875-4877) and the termination TGA codon (position 7848-7850) are underlined. The positions of intron 1 (7586-7822) and intron 2 (7942-8053) are indicated in bold italics.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: CPRO-DLO  
(B) STREET: Droevendaalsesteeg 1  
(C) CITY: Wageningen  
(D) STATE: Gelderland  
(E) COUNTRY: The Netherlands  
(F) POSTAL CODE (ZIP): Postbus 16 6700 AA

(A) NAME: Landbouw Universiteit Wageningen  
(B) STREET: Dreyenlaan 2  
(C) CITY: Wageningen  
(D) STATE: Gelderland  
(E) COUNTRY: Netherlands  
(F) POSTAL CODE (ZIP): Postbus 9101 6700 HB

(ii) TITLE OF INVENTION: Engineering nematode resistance in Solanaceae

(iii) NUMBER OF SEQUENCES: 7

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2739 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: *Gpa2* encoding sequence of *S. tuberosum*

## (ix) FEATURE:

(A) NAME/KEY: CDS  
(B) LOCATION: 1..2739

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

## 49

atg gct tat gct gct gtt act tcc ctt atg aga acc ata cat caa tca	48
Met Ala Tyr Ala Ala Val Thr Ser Leu Met Arg Thr Ile His Gln Ser	
1 5 10 15	
atg gaa ctt act gga tgt gat ttg caa ccg ttt tat gaa aag ctc aaa	96
Met Glu Leu Thr Gly Cys Asp Leu Gln Pro Phe Tyr Glu Lys Leu Lys	
20 25 30	
tct ttg aga gct att ctg gag aaa tcc tgc aat ata atg ggc gat cat	144
Ser Leu Arg Ala Ile Leu Glu Lys Ser Cys Asn Ile Met Gly Asp His	
35 40 45	
gag ggg tta aca atc ttg gaa gtt gaa atc ata gag gta gca tac aca	192
Glu Gly Leu Thr Ile Leu Glu Val Glu Ile Ile Glu Val Ala Tyr Thr	
50 55 60	
aca gaa gat atg gtt gac tcg gaa tca aga aat gtt ttt tta gca cgg	240
Thr Glu Asp Met Val Asp Ser Glu Ser Arg Asn Val Phe Leu Ala Arg	
65 70 75 80	
aat gtg ggg aaa aga agc agg gct atg tgg ggg att ttt ttc gtc ttg	288
Asn Val Gly Lys Arg Ser Arg Ala Met Trp Gly Ile Phe Phe Val Leu	
85 90 95	
gaa caa gca cta gaa tgc att gat tcc acc gtg aaa cag tgg atg gca	336
Glu Gln Ala Leu Glu Cys Ile Asp Ser Thr Val Lys Gln Trp Met Ala	
100 105 110	
aca tcg gac agc atg aaa gat cta aaa cca caa act agc tca ctt gtc	384
Thr Ser Asp Ser Met Lys Asp Leu Lys Pro Gln Thr Ser Ser Leu Val	
115 120 125	
agt tta cct gaa cat gat gtt gag cag ccc gag aat ata atg gtt ggc	432
Ser Leu Pro Glu His Asp Val Glu Gln Pro Glu Asn Ile Met Val Gly	
130 135 140	
cgt gaa aat gaa ttt gag atg atg ctg gat caa ctt gct aga gga gga	480
Arg Glu Asn Glu Phe Glu Met Met Leu Asp Gln Leu Ala Arg Gly Gly	
145 150 155 160	
agg gaa cta gaa gtt gtc tca atc gta ggg atg gga ggc atc ggg aaa	528
Arg Glu Leu Glu Val Val Ser Ile Val Gly Met Gly Gly Ile Gly Lys	
165 170 175	
aca act ttg gct gca aaa ctc tat agt gat cct tac att atg tct cga	576
Thr Thr Leu Ala Ala Lys Leu Tyr Ser Asp Pro Tyr Ile Met Ser Arg	
180 185 190	
ttt gat att cgt gca aaa gca act gtt tca caa gag tat tgt gtg aga	624
Phe Asp Ile Arg Ala Lys Ala Thr Val Ser Gln Glu Tyr Cys Val Arg	
195 200 205	
aat gta ctc cta ggc ctt ctt tct ttg aca agt gat gaa cct gat tat	672
Asn Val Leu Leu Gly Leu Leu Ser Leu Thr Ser Asp Glu Pro Asp Tyr	
210 215 220	

## 50

cag cta gcg gac caa ctg caa aag cat ctg aaa ggc agg aga tac ttg	720
Gln Leu Ala Asp Gln Leu Gln Lys His Leu Lys Gly Arg Arg Tyr Leu	
225 230 235 240	
gta gtc att gat gac ata tgg act aca gaa gct tgg gat gat ata aaa	768
Val Val Ile Asp Asp Ile Trp Thr Thr Glu Ala Trp Asp Asp Ile Lys	
245 250 255	
cta tgt ttc cca gac tgc gat aat gga agc aga ata ctc ctg act act	816
Leu Cys Phe Pro Asp Cys Asp Asn Gly Ser Arg Ile Leu Leu Thr Thr	
260 265 270	
cgg aat gtg gaa gtg gct gaa tat gct agc tca ggt aag cct cct cat	864
Arg Asn Val Glu Val Ala Glu Tyr Ala Ser Ser Gly Lys Pro Pro His	
275 280 285	
cac atg cgc ctc atg aat ttt gac gaa agt tgg aat tta cta cac aaa	912
His Met Arg Leu Met Asn Phe Asp Glu Ser Trp Asn Leu Leu His Lys	
290 295 300	
aag atc ttt gaa aaa gaa ggt tct tat tct cct gaa ttt gaa aat att	960
Lys Ile Phe Glu Lys Glu Gly Ser Tyr Ser Pro Glu Phe Glu Asn Ile	
305 310 315 320	
ggg aaa caa att gca tta aaa tgt gga ggg tta cct cta gca att act	1008
Gly Lys Gln Ile Ala Leu Lys Cys Gly Gly Leu Pro Leu Ala Ile Thr	
325 330 335	
ttg att gct gga ctt ctc tcc aaa atc agt aaa aca ttg gat gag tgg	1056
Leu Ile Ala Gly Leu Leu Ser Lys Ile Ser Lys Thr Leu Asp Glu Trp	
340 345 350	
caa aat gtt gcg gag aat gta cgt tcg gtg gta agc aca gat ctt gaa	1104
Gln Asn Val Ala Glu Asn Val Arg Ser Val Val Ser Thr Asp Leu Glu	
355 360 365	
gca aaa tgc atg aga gtg ttg gct ttg agt tac cat cac ttg cct tct	1152
Ala Lys Cys Met Arg Val Leu Ala Leu Ser Tyr His His Leu Pro Ser	
370 375 380	
cac cta aaa ccg tgt ttt ctg tat ttt gca att ttc gca gag gat gaa	1200
His Leu Lys Pro Cys Phe Leu Tyr Phe Ala Ile Phe Ala Glu Asp Glu	
385 390 395 400	
cgg att tat gta aat aaa ctt gtt gag tta tgg gcc gta gag ggg ttt	1248
Arg Ile Tyr Val Asn Lys Leu Val Glu Leu Trp Ala Val Glu Gly Phe	
405 410 415	
ttg aat gaa gaa gag gga aaa agc ata gaa gag gtg gca gaa aca tgt	1296
Leu Asn Glu Glu Glu Gly Lys Ser Ile Glu Glu Val Ala Glu Thr Cys	
420 425 430	
ata aac gaa ctt gta gat aga agt cta att tct atc cac aat gtg agt	1344
Ile Asn Glu Leu Val Asp Arg Ser Leu Ile Ser Ile His Asn Val Ser	
435 440 445	



## 51

ttt gat ggg gaa aca cag aga tgt gga atg cat gat gtg acc cgt gaa	1392
Phe Asp Gly Glu Thr Gln Arg Cys Gly Met His Asp Val Thr Arg Glu	
450 455 460	
ctc tgt ttg agg gaa gct cga aac atg aat ttt gtg aat gtt atc aga	1440
Leu Cys Leu Arg Glu Ala Arg Asn Met Asn Phe Val Asn Val Ile Arg	
465 470 475 480	
gga aag agt gat caa aat tca tgt gca caa tcc atg cag tgt tcc ttt	1488
Gly Lys Ser Asp Gln Asn Ser Cys Ala Gln Ser Met Gln Cys Ser Phe	
485 490 495	
aag agt cga agt cgg atc agt atc cat aat gag gaa gaa ttg gtt tgg	1536
Lys Ser Arg Ser Arg Ile Ser Ile His Asn Glu Glu Glu Leu Val Trp	
500 505 510	
tgt cgt aac agc gag gct cat tct atc atc acg ttg tgt ata ttc aaa	1584
Cys Arg Asn Ser Glu Ala His Ser Ile Ile Thr Leu Cys Ile Phe Lys	
515 520 525	
tgc gtc aca ctg gaa ttg tct ttc aag cta gta aga gta cta gat ctt	1632
Cys Val Thr Leu Glu Leu Ser Phe Lys Leu Val Arg Val Leu Asp Leu	
530 535 540	
ggg ttg act aca tgc cca att ttt ccc agt gga gta ctt tct cta att	1680
Gly Leu Thr Thr Cys Pro Ile Phe Pro Ser Gly Val Leu Ser Leu Ile	
545 550 555 560	
cat ttg aga tac cta tct ttg cgt ttt aat cct cgc tta cag cag tat	1728
His Leu Arg Tyr Leu Ser Leu Arg Phe Asn Pro Arg Leu Gln Gln Tyr	
565 570 575	
cga gga tcg aaa gaa gct gtt ccc tca tca ata ata gac att cct cta	1776
Arg Gly Ser Lys Glu Ala Val Pro Ser Ser Ile Ile Asp Ile Pro Leu	
580 585 590	
tcg ata tca agc cta tgc tat ctg caa act ttt aaa ctt tac cat cca	1824
Ser Ile Ser Ser Leu Cys Tyr Leu Gln Thr Phe Lys Leu Tyr His Pro	
595 600 605	
ttt ccc aat tgt tat cct ttc ata tta cca tcg gaa att ttg aca atg	1872
Phe Pro Asn Cys Tyr Pro Phe Ile Leu Pro Ser Glu Ile Leu Thr Met	
610 615 620	
cca caa ttg agg aag ctg tgt atg ggc tgg aat tac ttg cgg agt cat	1920
Pro Gln Leu Arg Lys Leu Cys Met Gly Trp Asn Tyr Leu Arg Ser His	
625 630 635 640	
gag cct aca gag aac aga ttg gtt ttg aaa agt ttg caa tgc ctc aat	1968
Glu Pro Thr Glu Asn Arg Leu Val Leu Lys Ser Leu Gln Cys Leu Asn	
645 650 655	
gaa ttg aat cct cgg tat tgt aca ggg tct ttt tta aga cta ttt ccc	2016
Glu Leu Asn Pro Arg Tyr Cys Thr Gly Ser Phe Leu Arg Leu Phe Pro	
660 665 670	

52

aat tta aag aag ttg gaa gta ttt ggc gtc aaa gag gac ttt cgc aat	2064
Asn Leu Lys Lys Leu Glu Val Phe Gly Val Lys Glu Asp Phe Arg Asn	
675 680 685	
cac aag gac ctg tat gat ttt cgc tac tta tat cag ctc gag aaa ttg	2112
His Lys Asp Leu Tyr Asp Phe Arg Tyr Leu Tyr Gln Leu Glu Lys Leu	
690 695 700	
gca ttt agt act tat tat tca tct tct gct tgc ttt cta aaa aac act	2160
Ala Phe Ser Thr Tyr Tyr Ser Ser Ser Ala Cys Phe Leu Lys Asn Thr	
705 710 715 720	
gca cct tta ggt tct act ccg caa gat cct ctg agg ttt cag atg gaa	2208
Ala Pro Leu Gly Ser Thr Pro Gln Asp Pro Leu Arg Phe Gln Met Glu	
725 730 735	
aca ttg cac tta gag act cat tcc agg gca act gca cct cca act gat	2256
Thr Leu His Leu Glu Thr His Ser Arg Ala Thr Ala Pro Pro Thr Asp	
740 745 750	
gtt cca act ttc ctc tta cct cct ccg gat tgt ttt cca caa aac ctt	2304
Val Pro Thr Phe Leu Leu Pro Pro Pro Asp Cys Phe Pro Gln Asn Leu	
755 760 765	
aag agt tta act ttt agc gga gat ttc ttt ttg gca tgg aag gat ttg	2352
Lys Ser Leu Thr Phe Ser Gly Asp Phe Phe Leu Ala Trp Lys Asp Leu	
770 775 780	
agc att gtt ggt aaa tta ccc aaa ctc gag gtc ctt caa cta tca cac	2400
Ser Ile Val Gly Lys Leu Pro Lys Leu Glu Val Leu Gln Leu Ser His	
785 790 795 800	
aat gcc ttc aaa ggc gag gag tgg gaa gta gtt gag gaa ggg ttt cct	2448
Asn Ala Phe Lys Gly Glu Glu Trp Glu Val Val Glu Glu Gly Phe Pro	
805 810 815	
cac ttg aag ttc ttg ttt ctg gat agc ata tac att cgg tac tgg aga	2496
His Leu Lys Phe Leu Phe Leu Asp Ser Ile Tyr Ile Arg Tyr Trp Arg	
820 825 830	
gct agt agt gat cac ttt cca tac ctt gaa cga ctt ttt ctt agc gat	2544
Ala Ser Ser Asp His Phe Pro Tyr Leu Glu Arg Leu Phe Leu Ser Asp	
835 840 845	
tgc ttt tat ttg gat tca atc cct cga gat ttt gca gat ata acc aca	2592
Cys Phe Tyr Leu Asp Ser Ile Pro Arg Asp Phe Ala Asp Ile Thr Thr	
850 855 860	
cta gct ctt att gat ata ttt cgc tgc caa caa tct gtt ggg aat tcc	2640
Leu Ala Leu Ile Asp Ile Phe Arg Cys Gln Gln Ser Val Gly Asn Ser	
865 870 875 880	
gcc aag caa att caa cag gac att caa gac aac tat gga agc tct atc	2688
Ala Lys Gln Ile Gln Gln Asp Ile Gln Asp Asn Tyr Gly Ser Ser Ile	
885 890 895	

gag gtc cat act cgt tat ctt tat cga aat gga gca ttt ttg gta gtg 2736  
 Glu Val His Thr Arg Tyr Leu Tyr Arg Asn Gly Ala Phe Leu Val Val  
                   900                                  905                                  910

tga 2739  
 \*

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2976 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Gpa2* coding and non coding sequence of *S. tuberosum*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

ATGGCTTATG CTGCTGTTAC TTCCCTTATG AGAACCATAC ATCAATCAAT GGAACTTACT	60
GGATGTGATT TGCAACCGTT TTATGAAAAG CTCAAATCTT TGAGAGCTAT TCTGGAGAAA	120
TCCTGCAATA TAATGGGCGA TCATGAGGGG TTAACAATCT TGGAAGTTGA AATCATAGAG	180
GTAGCATACA CAACAGAAGA TATGGTTGAC TCGGAATCAA GAAATGTTTT TTTAGCACGG	240
AATGTGGGGA AAAGAAGCAG GGCTATGTGG GGGATTTTTT TCGTCTTGGA ACAAGCACTA	300
GAATGCATTG ATTCCACCGT GAAACAGTGG ATGGCAACAT CGGACAGCAT GAAAGATCTA	360
AAACCACAAA CTAGCTCACT TGTCAGTTTA CCTGAACATG ATGTTGAGCA GCCCCGAGAAT	420
ATAATGGTTG GCCGTGAAAA TGAATTTGAG ATGATGCTGG ATCAACTTGC TAGAGGAGGA	480
AGGGAAGTAG AAGTTGTCTC AATCGTAGGG ATGGGAGGCA TCGGGAAAAC AACTTTGGCT	540
GCAAAACTCT ATAGTGATCC TTACATTATG TCTCGATTTG ATATTCGTGC AAAAGCAACT	600
GTTTCACAAG AGTATTGTGT GAGAAATGTA CTCCTAGGCC TTCTTTCTTT GACAAGTGAT	660
GAACCTGATT ATCAGCTAGC GGACCAACTG CAAAAGCATC TGAAAGGCAG GAGATACTTG	720
GTAGTCATTG ATGACATATG GACTACAGAA GCTTGGGATG ATATAAAACT ATGTTTCCCA	780
GACTGCGATA ATGGAAGCAG AATACTCCTG ACTACTCGGA ATGTGGAAGT GGCTGAATAT	840
GCTAGCTCAG GTAAGCCTCC TCATCACATG CGCCTCATGA ATTTTGACGA AAGTTGGAAT	900

TTACTACACA AAAAGATCTT TGAAAAAGAA GGTTCCTTATT CTCCTGAATT TGAAAATATT	960
GGGAAACAAA TTGCATTAAA ATGTGGAGGG TTACCTCTAG CAATTACTTT GATTGCTGGA	1020
CTTCTCTCCA AAATCAGTAA AACATTGGAT GAGTGGCAAA ATGTTGCGGA GAATGTACGT	1080
TCGGTGGTAA GCACAGATCT TGAAGCAAAA TGCATGAGAG TGTGGCTTT GAGTTACCAT	1140
CACTTGCCCTT CTCACCTAAA ACCGTGTTTT CTGTATTTTG CAATTTTCGC AGAGGATGAA	1200
CGGATTTTATG TAAATAAACT TGTTGAGTTA TGGGCCGTAG AGGGGTTTTT GAATGAAGAA	1260
GAGGGAAAAA GCATAGAAGA GGTGGCAGAA ACATGTATAA ACGAACTTGT AGATAGAAGT	1320
CTAATTTCTA TCCACAATGT GAGTTTTGAT GGGGAAACAC AGAGATGTGG AATGCATGAT	1380
GTGACCCGTG AACTCTGTTT GAGGGAAGCT CGAAACATGA ATTTTGTGAA TGTATCAGA	1440
GGAAAGAGTG ATCAAAATTC ATGTGCACAA TCCATGCAGT GTTCCTTTAA GAGTCGAAGT	1500
CGGATCAGTA TCCATAATGA GGAAGAATTG GTTTGGTGTC GTAACAGCGA GGCTCATTCT	1560
ATCATCACGT TGTGTATATT CAAATGCGTC ACACTGGAAT TGTCTTTCAA GCTAGTAAGA	1620
GTACTAGATC TTGGTTTGAC TACATGCCCA ATTTTCCCA GTGGAGTACT TTCTCTAATT	1680
CATTTGAGAT ACCTATCTTT GCGTTTTAAT CCTCGCTTAC AGCAGTATCG AGGATCGAAA	1740
GAAGCTGTTT CCTCATCAAT AATAGACATT CCTCTATCGA TATCAAGCCT ATGCTATCTG	1800
CAAACTTTTA AACTTTACCA TCCATTTCCC AATTGTTATC CTTTCATATT ACCATCGGAA	1860
ATTTTGACAA TGCCACAATT GAGGAAGCTG TGTATGGGCT GGAATTACTT GCGGAGTCAT	1920
GAGCCTACAG AGAACAGATT GGTTTTGAAA AGTTTGCAAT GCCTCAATGA ATTGAATCCT	1980
CGGTATTGTA CAGGGTCTTT TTTAAGACTA TTTCCCAATT TAAAGAAGTT GGAAGTATTT	2040
GGCGTCAAAG AGGACTTTCG CAATCACAAG GACCTGTATG ATTTTCGCTA CTTATATCAG	2100
CTCGAGAAAT TGGCATTTAG TACTTATTAT TCATCTTCTG CTTGCTTTCT AAAAAACACT	2160
GCACCTTTAG GTTCTACTCC GCAAGATCCT CTGAGGTTTC AGATGGAAAC ATTGCACTTA	2220
GAGACTCATT CCAGGGCAAC TGCACCTCCA ACTGATGTTT CAACTTTCCT CTTACCTCCT	2280
CCGGATTGTT TTCCACAAAA CCTTAAGAGT TTAACTTTTA GCGGAGATTT CTTTTTGGCA	2340
TGGAAGGATT TGAGCATTGT TGGTAAATTA CCCAACTCG AGGTCCTTCA ACTATCACAC	2400
AATGCCTTCA AAGGCGAGGA GTGGGAAGTA GTTGAGGAAG GGTTCCTCA CTTGAAGTTC	2460
TTGTTTCTGG ATAGCATATA CATTCGGTAC TGGAGAGCTA GTAGTGATCA CTTTCCATAC	2520
CTTGAACGAC TTTTCTTAG CGATTGCTTT TATTTGGATT CAATCCCTCG AGATTTTGCA	2580

GATATAACCA CACTAGCTCT TATTGATATA TTTCGCTGCC AACAACTCTGT TGGGAATTCC 2640  
 GCCAAGCAAA TTCAACAGGA CATTCAAGAC AACTATGGAA GCTCTATCGA GGTCCATACT 2700  
 CGTTATCTTT **AGTAAGACAT CTTCTTCCTT GATTTACAAC AATATTTAAC TCATCATCAT** 2760  
**AGTAAACTCG ATAATAATCT GGATAATAGC TTTAGTAAGT CAAATTGCAC CAATTCAACA** 2820  
**AAAGTTCTTG ATGCTGTCAT TGTGATTGAT TCGAATCCTT CCAATATTGT GTAACCTGTT** 2880  
**ATACTTGCAAT GTTCATTCTT GATTTTGGGA AGTGTAACAT TTCCATTTTT CATCTTGATT** 2940  
**TTGGGAAG**TC GAAATGGAGC ATTTTGGTA GTGTGA 2976

## (2) INFORMATION FOR SEQ ID NO: 3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10329 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: *XbaI-XbaI* pBINRGH2 fragment containing *Gpa2* promoter, coding and non coding sequence of *S. tuberosum*.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CTAGAGATTG GAATGGAGTG ATTCTTAGGG GTTCTTTTTT GAATTAATAT GAGGGTTAGT 60  
 ATTCAATCTT CAATTCGACA TTTTCTCATA ATTTCTTTAT CTGTTTATTT TTCCTATTCG 120  
 TAAATCTCTT GGGAAAAATT GGGGTTTTAT CGATTGGGAC TCCTTTTTGA TGAAAAAGGT 180  
 ATATTTACGA TCTTTATGTT ATGGGTAAAC TGATTTTAAC ATAAAATTAT TGATTCATCG 240  
 ATTATTTTTA TCATATTAAC CGCGTACAAT TTGGACTTTC CCGGTAAAGT TAAAGTATGA 300  
 TAAATTGAGA ATTTCAAGGT CGATCTTAGC TCCATTTTTG ATGAAATTTT ATATTTGAAC 360  
 TTATCTAAGC ATGGGTAAGA TGTTTTTCAA GAAATATTTT ATTTTCGAGT CGGGGTTTTG 420  
 GATTCGAATA TTTTAGGCTT CTTCAAGAAT GTAGATTTTT GTTTAAATTG AGTTTGTGAA 480  
 TTGATTTCAA CTCCATTTTC AAATTGGTTT TCACCATTAG CTTCCAAATA CTTTAAGGAT 540  
 CATTTTACAT CAAAAAATTC CAGATTGGG TATCGTTTTT CCGTATGAGA CTTTTGGACC 600

GTTTTGCCCC	TTTTCCCTAA	ATTTCTTGAT	TTTGGTGTC	TTGGACTCGA	ATTGTGATTG	660
TGAATAATTG	TTTGAATAGA	TTATCGTGAT	CCAGATTATA	CTTGGAAGG	AAAGGCTCAA	720
GTCAAGTAAC	TTTTGGAGTT	CGTTTTAAGG	CAAGTGGCTT	CCAAACTTTG	TAAAACTCTT	780
AGACTACGCA	TGACTACTTT	CCTAATTATG	TTGGGGAGTA	ATGGGGGATT	GAGGATGGGT	840
TTTATTTGTT	GATTGAAATT	GTTGTAAATG	AAAGATGGGG	AATAAAACGA	GCTAAATGTG	900
TTATGTGTGA	CTTGAATTTG	TTTGAATAAG	TCATGTGATA	ACTGATATTG	AGGGATAGAA	960
GAGCATGAGC	AGGCTATGAT	TGATACAGAC	ATTGATGTTG	AGGCAGATGA	TGTGTAATAC	1020
TATGATGTGG	TCGTGATATG	GTTGTGATTG	AGACATGTGA	TGTGTAATAC	TATGATGTGG	1080
TCGTGATATG	GTTGTGATTG	AGACAGGTGA	TGTGTAATAC	TATGATGTGG	TCGTGATATG	1140
GTTGTGATTG	AGACAGGTGA	TGTGTAATAC	TATGATGTGG	TCGTGATATG	GTTGTGACTG	1200
AGACAGGTGA	TGTGTAATAC	TATGATGTGG	TCGTGATATG	GTTGTGATTG	AGACAGATGA	1260
TGTGTAATAC	GATGATGTGA	TCGTGATATG	ATTGTGATTG	ATTACATGTG	CATATTCATT	1320
ATTCATCCCA	TGTGTGAACT	ATCTGTTGCA	TGAGTTCTGA	GACACTGATA	TGAGGATGGA	1380
TGGATATGAG	ACACAGTTGA	GACTAGCTCC	GGCTAGAGAT	GTATGAGATG	GACTAGCTCC	1440
GGCTAGCGAT	TTGGATGCCG	ATGGGATCTG	GTTCCGGCGG	TGATACATGG	TCCATGTGTG	1500
GCCCCATGG	GTTCTGATTT	GAGTATTCAA	CGCGGACTGA	TTACGTCAAC	AGATGTGTAT	1560
CGTAGGACAG	ACATGTATCA	CGACTACATG	ACATCATTAT	TGCATTTTGC	ATCGCATTTG	1620
CCTTATCTTT	GTCTGTGATG	TGTGGATTGT	ATCGGTTTAC	CCTTTTTTATG	TGGAATTTGA	1680
TCTACTTGCT	CTTATTTGTT	GATCTGAGGT	TGATGAGGAT	ATACTGTTGG	TTCTGGCTGT	1740
TGAATATGAT	CTGTTTAGTA	TAGGTTGGTT	GGTTTGCTGC	TAGATTGAAG	TTTCGGTGGT	1800
TCGGTTGGGA	TTGAAAGGAG	TTGTTTGTAG	CTGCTAGTTT	TGCTTAGTTT	AGAGTTACTT	1860
GCGAGTACCT	GTGGTTTTTCG	GTA CTCACCC	TTGCTTCTAC	ACAATTGTGT	AGGTTGACAG	1920
CTCTCTCTCA	GATATTTTCT	TTAGCAGATT	GAGCTTTGAG	ACATACTCGA	GAGGTAGCGG	1980
TTCATTCCAG	ACGTGCCCTT	GAGTTATCTT	TACTTTCAGT	TTTGTTCTAT	TCGAGAACTA	2040
TACTCTGAGA	CTTGATATAT	TTTATTCGAA	TTCTGTATTT	AGAGGTTTGT	ACATGTGACA	2100
ACCAAATTCT	GGGTAGTGTT	AAGTCTTAAT	TAAAGTTTTT	TGCTTATTTA	TTATCTTTTA	2160
TTCTCGTATT	TCTACTTCTC	TATCGTTGTG	GTTGGGTTAG	GCTGACGTGT	CTGGTGGGAA	2220
ACGGACATGT	GCCATCACAT	CCGGATTTGG	GGTGTGACAA	ATATTTTGTT	AGTTATATAC	2280

AAAATTGTAT	GTAGTATATG	TATATTTTCT	GCTTTCATCA	CAATTGTATA	TAGATATTTG	2340
TATATTTTGT	TAGTTATATA	CAAAATTGCT	TGAAGTATAT	GTATATTTTC	TGCTTAAATC	2400
ATAATTGTAT	ATATATATAT	ATATATATAT	ATTTCTATAT	TTTGTAAGTT	ATATACAATA	2460
GTATGAATTA	AACAATATAC	AAACCTTACA	TTATTATATA	TACAGTTAGG	TTACACCAAA	2520
AATTATCAAA	TTAAAGCACA	ACTTTTTTAT	CGAATCATAT	ACAATTCATA	TATATAATTG	2580
ACTTAGTAAT	TTTATACAAC	TACTTACACT	TCTACATGGT	ATAAGAATTT	TGCACAATTA	2640
CTTACATATA	TACAATATTA	TCAATTAAAC	AATATACAAA	TCGTATAACT	TATATATACA	2700
GTAAATTAC	AACAACAACA	ACAAAAATTA	TCAAATTAAA	GCACACCGTT	GTTGTCGAAT	2760
CATATACACT	CCATATATAC	AAATTGTGTC	ATTCAATTTT	TCGAACAAAA	AATTAGAATT	2820
GAATTGTAA	TATAAAATTT	ATCTAATATT	GTATAAACAA	AATTAAATTA	TTGCAAACCA	2880
TTAGAATGAA	AAAAACAAAA	ATAAACCGTT	TTCCAAAATT	TCAATTATAT	ACTATACAAA	2940
TCAATTGTAT	ACTTTCTTGC	CGTTCAAAAC	ATGAAGTTTC	CTTGAAAGAA	ACGCTTACCT	3000
AGCGTTGAAT	ATACAAGAAT	ATTGATTAAT	CGTATGCTTC	AGTCGTTTGA	GGAACCCAGT	3060
TGTTATTGTG	TTTCTATTGC	TATAGAACTC	CTTTTTGGAA	AAATATTTGA	TTTTGGACGA	3120
TTAGCTTGAA	TCATGGGATT	ATATAAAATT	TTTATTACCG	TATTTAGCAC	TCATGTATCC	3180
ATTTATTAAA	AAAAAATTGT	ATAAATTATA	TTTTTAAAAG	AAAATATACA	AAATTAATGC	3240
TTCATAGCAA	ACTAACTAT	ACCCATTGAA	TGTAATTACT	AACTATACC	TATAGAGCGT	3300
TATTTCATTA	AATACGTTTA	TCATATATGA	AGTTTTCCCT	CAAGAGATCC	TACACCTTAT	3360
ATATAGCTTC	TCAAATGTGG	AAATTCAATC	TCACACCCAA	CAATCTTTCC	CTCAGACTAA	3420
GTTTCATGGC	CCAATATCAC	AATGATCCAC	GAGTCAATTC	ATGAGATTCA	CTATGTGTGT	3480
CACCCACATC	GTCTAAGTAT	TTTATGGCAA	TCAAGCCCTA	CAACTTGCTT	CTTCTTTATA	3540
TATATATATA	TATATATATA	TATATATATA	TATATATGTG	TGTGTGTGTG	TGTGTGTGTG	3600
CGCATCTCTA	ATTAATCTCG	TAAAGGGATT	AAGGGGCCAA	TTTCAAAGAA	TTAGGCGATT	3660
TTCTTAGTTT	TTCGTGTGTG	TTAACCCTA	GGTATTTTGG	TGATATGGTT	TTCGGATGAT	3720
TTATTTTGTG	CAACTTATAT	GGAACCCTTC	GTAGGGAGTT	AGTCTCACAC	TTTTTAGAGT	3780
CCATTTTGGG	CATTCAGGGG	CTAATTTATA	GGAAATAGGT	GATCTTCTCA	GTTTGTCTGT	3840
ATTAGCCCAT	GAATATTTTG	GTGATATGTC	TTCCGAATAA	TTTCTTTGTA	AAATCTTTAC	3900
GGGACCCTCC	ATAGGGAGTT	AGTGGAGCAG	TACGTATAGT	CTCACAATTT	TAGAGTTCAT	3960

TTTGGGCATT TAGGGGCCAA TTTACAGGAT TTAGGCGACT TTCTCAGTGT TTTGTGTGTG	4020
TTAGCCCATT AATAGTTGGT GATATGACTT TCAGACGATT TCTTTGCTAC ACATTTACGG	4080
AACCCTCTGT AGGAAGTCGG GGGAGCAATA CGTACAATCT CACAATTTTA GAGTCCATTT	4140
TAGGCATTTA GGGGCCAATT TAAAGAAATT GGACAATTTT CTCAGTTTTT CGTGTCTGTT	4200
AGCCATTAAT ATATTGGTGA ATATGACCTA CAGATGATTT CTAATCGAAA TCTTTACGAA	4260
ACCCTCAGTA GGGAGTTGGG GGAGCAATAC GTACCGTCTG ACAATTTTTTA GAGTCCATTT	4320
TGGGCATTTA AGGGCCAATT TACAGGAATT AGACGATTTT CTTAGTATTT TTTTCATGTGT	4380
TAGCCCATAA ATATTTTGTT GATTTGACTT TTAGAGTCTA AACTTCTCAT GTATATTAAG	4440
AGATATTTAT GCTTGTTAA TTGAATCGAA CTAGGAATAG AGAAATTCCT ACTTGGATCT	4500
TAATATTTCT CTCTCTTTGA TTTGGAAAAT TCTAGGAAGT TGCTTTCAAT GGAATTAAAA	4560
TCATCAATCT CTTGTATGTA AGAAACATAC TTATATTCAT GAATAGATAT GTTTAGGGTC	4620
TAATAATGAA TTATCACAAT TTTTCTACT TTTTCTTGTC AGAGTCCTGC CTTTTCTTTT	4680
TTCTTTTTTA ACTTTGGTCT CTGCTTTTGT CTACATGATG ATAAGGTTGG TGGACCTAGC	4740
TGGAAATGTG ATGGAAATAG CTAGTAAAAG AAAGAACTTT GCATTTTCTG TTTTCTTAAA	4800
AACTGATAAA TTACATAACT TGTGGCAATT TGTCCATTTT CATACTGAGA GATATTTCTA	4860
TTTTTTTTTG ATAT <u>ATG</u> GCT TATGCTGCTG TTA <del>CT</del> CCCT TATGAGAACC ATACATCAAT	4920
CAATGGAACT TACTGGATGT GATTTGCAAC CGTTTTATGA AAAGCTCAA TCTTTGAGAG	4980
CTATTCTGGA GAAATCCTGC AATATAATGG GCGATCATGA GGGGTTAACA ATCTTGGAAG	5040
TTGAAATCAT AGAGGTAGCA TACACAACAG AAGATATGGT TGA <del>CT</del> CGGAA TCAAGAAATG	5100
TTTTTTTAGC ACGGAATGTG GGGAAAAGAA GCAGGGCTAT GTGGGGGATT TTTTTCGTCT	5160
TGGAACAAGC ACTAGAATGC ATTGATTCCA CCGTGAAACA GTGGATGGCA ACATCGGACA	5220
GCATGAAAGA TCTAAAACCA CAACTAGCT CACTTGTCAG TTTACCTGAA CATGATGTTG	5280
AGCAGCCCGA GAATATAATG GTTGGCCGTG AAAATGAATT TGAGATGATG CTGGATCAAC	5340
TTGCTAGAGG AGGAAGGGAA CTAGAAGTTG TCTCAATCGT AGGGATGGGA GGCATCGGGA	5400
AAACAACTTT GGCTGCAAAA CTCTATAGTG ATCCTTACAT TATGTCTCGA TTTGATATTC	5460
GTGCAAAAGC AACTGTTTCA CAAGAGTATT GTGTGAGAAA TGTACTCCTA GGCCTTCTTT	5520
CTTTGACAAG TGATGAACCT GATTATCAGC TAGCGGACCA ACTGCAAAAAG CATCTGAAAG	5580
GCAGGAGATA CTTGGTAGTC ATTGATGACA TATGGACTAC AGAAGCTTGG GATGATATAA	5640



AACATATGTTT	CCCAGACTGC	GATAATGGAA	GCAGAATACT	CCTGACTACT	CGGAATGTGG	5700
AAGTGGCTGA	ATATGCTAGC	TCAGGTAAGC	CTCCTCATCA	CATGCGCCTC	ATGAATTTTG	5760
ACGAAAGTTG	GAATTTACTA	CACAAAAAGA	TCTTTGAAAA	AGAAGGTTCT	TATTCTCCTG	5820
AATTTGAAAA	TATTGGGAAA	CAAATTGCAT	TAAAATGTGG	AGGGTTACCT	CTAGCAATTA	5880
CTTTGATTGC	TGGACTTCTC	TCCAAAATCA	GTAAAACATT	GGATGAGTGG	CAAAATGTTG	5940
CGGAGAATGT	ACGTTCCGGT	GTAAGCACAG	ATCTTGAAGC	AAAATGCATG	AGAGTGTGG	6000
CTTTGAGTTA	CCATCACTTG	CCTTCTCACC	TAAAACCGTG	TTTTCTGTAT	TTTGCAATTT	6060
TCGCAGAGGA	TGAACGGATT	TATGTAAATA	AACCTGTTGA	GTTATGGGCC	GTAGAGGGGT	6120
TTTTGAATGA	AGAAGAGGGA	AAAAGCATAG	AAGAGGTGGC	AGAAACATGT	ATAAACGAAC	6180
TTGTAGATAG	AAGTCTAATT	TCTATCCACA	ATGTGAGTTT	TGATGGGGAA	ACACAGAGAT	6240
GTGGAATGCA	TGATGTGACC	CGTGAACCTC	GTTTGAGGGA	AGCTCGAAAC	ATGAATTTTG	6300
TGAATGTTAT	CAGAGGAAAG	AGTGATCAAA	ATTCATGTGC	ACAATCCATG	CAGTGTTCCT	6360
TTAAGAGTCG	AAGTCGGATC	AGTATCCATA	ATGAGGAAGA	ATTGGTTTGG	TGTCGTAACA	6420
GCGAGGCTCA	TTCTATCATC	ACGTTGTGTA	TATTCAAATG	CGTCACACTG	GAATTGTCTT	6480
TCAAGCTAGT	AAGAGTACTA	GATCTTGGTT	TGACTACATG	CCCAATTTTT	CCCAGTGGAG	6540
TACTTTCTCT	AATTCATTTG	AGATACCTAT	CTTTGCGTTT	TAATCCTCGC	TTACAGCAGT	6600
ATCGAGGATC	GAAAGAAGCT	GTTCCCTCAT	CAATAATAGA	CATTCCTCTA	TCGATATCAA	6660
GCCTATGCTA	TCTGCAAAC	TTTAAACTTT	ACCATCCATT	TCCAATTGT	TATCCTTTCA	6720
TATTACCATC	GGAAATTTTG	ACAATGCCAC	AATTGAGGAA	GCTGTGTATG	GGCTGGAATT	6780
ACTTGCGGAG	TCATGAGCCT	ACAGAGAACA	GATTGGTTTT	GAAAAGTTTG	CAATGCCTCA	6840
ATGAATTGAA	TCCTCGGTAT	TGTACAGGGT	CTTTTTTAAG	ACTATTTCCC	AATTTAAAGA	6900
AGTTGGAAGT	ATTTGGCGTC	AAAGAGGACT	TTCGCAATCA	CAAGGACCTG	TATGATTTTC	6960
GCTACTTATA	TCAGCTCGAG	AAATTGGCAT	TTAGTACTTA	TTATTCATCT	TCTGCTTGCT	7020
TTCTAAAAAA	CACCTGCACCT	TTAGGTTCTA	CTCCGCAAGA	TCCTCTGAGG	TTTCAGATGG	7080
AAACATTGCA	CTTAGAGACT	CATTCCAGGG	CAACTGCACC	TCCAACATGAT	GTTCCAACCT	7140
TCCTCTTACC	TCCTCCGGAT	TGTTTTCCAC	AAAACCTTAA	GAGTTTAACT	TTTAGCGGAG	7200
ATTTCTTTTT	GGCATGGAAG	GATTTGAGCA	TTGTTGGTAA	ATTACCCAAA	CTCGAGGTCC	7260
TTCAACTATC	ACACAATGCC	TTCAAAGGCG	AGGAGTGGGA	AGTAGTTGAG	GAAGGGTTTC	7320

CTCACTTGAA GTTCTTGTTT CTGGATAGCA TATACATTCG GTACTGGAGA GCTAGTAGTG	7380
ATCACTTTCC ATACCTTGAA CGACTTTTTTC TTAGCGATTG CTTTTATTTG GATTCAATCC	7440
CTCGAGATTT TGCAGATATA ACCACACTAG CTCTTATTGA TATATTTTCGC TGCCAACAAT	7500
CTGTTGGGAA TTCCGCCAAG CAAATTCAAC AGGACATTCA AGACAACTAT GGAAGCTCTA	7560
TCGAGGTCCA TACTCGTTAT CTTTAGTAAG ACATCTTCTT CTTTGATTTA CAACAATATT	7620
TAACTCATCA TCATAGTAAA CTCGATAATA ATCTGGATAA TAGCTTTAGT AAGTCAAATT	7680
GCACCAATTC AACAAAAGTT CTTGATGCTG TCATTGTGAT TGATTGGAAT CTTTCCAATA	7740
TTGTGTAACT TGTTATACTT GCATGTTTCT TCTTGATTTT GGAAGTGTA ACATTTCCAT	7800
TTTTTCATCTT GATTTTGGGA AGTCGAAATG GAGCATTTTT GGTAGTGGA CAACAGATGA	7860
AGATGATGAT GATAGTGTGA CAACAGATGA AGATGAAGAT GAAGACTTTG AGAAAGAAGT	7920
TGCTTCTTGC GGCAATAATG TGTAAAGTTCT TATACCTGCA TGCTCATTCT TGCTATAATG	7980
TTCTCTTGTT CCTTAATTAT GGGACATCTA ACATATTATT TTCCATTTTT TGCTCTTTT	8040
TTTTTTCCTG CAGCGTGTAG TTAAGGTGTT CTGAGGACTA GCCAGTTCTC TGAAATAAAT	8100
GTCAAATCAG AAGCCAAATG TGTGAGTGTG TGTTTTGTTT GTTTTCATTT TTTCTGCATA	8160
AGGTGGCAGG ATGATTGCAA ATGGCTTGTA ATTTAATTGT ATATGATATT TCGTATAGCC	8220
ATTTGCCAGT GGTTTTTTTAG ATACTCCAAA TTTTATGTAC ATACATAATG GTATAGGCCA	8280
GAACAGGCTC CATATATAAC GTGTGTTTCC TTTCTTGGA GTCTCAATC TACCTCGCAA	8340
AGGAAGACAG ACGGCTAAAT CAAGAAAGAA ATTTTTTTGA AAATCATGTG GCTAGTTGTT	8400
CAACTTTATA CAAGTTTATG TGCATACTTG TGCATACCCA AAGTTGAATA ACATAAACAT	8460
AAAATGAAGT CAAGTTAAAT GGCACATTTA TGTATTATGC CTTTGAATT TCATTAATAG	8520
TGAAAAATCCT GAATCATATT CAGATTCCAT CACTAATCGT TGAACCATGT TAATTTACTA	8580
TGTATTATCT AATGGATTTT TTTGCTATCT TATTTATAAT TGTTCAAAGT TTTGTTAATT	8640
ATCTTTAGCA TAATATCTGA TTATATTATT TTGATATACT TTCTCTATCC CTAATTACTT	8700
GTCCATTTTT GAATTGGCAC ACCTATTAAG AAAATAATTA TTGAAATAGT GAGTTTACCA	8760
TTTTACCCAT ATTAATTATG AAGTGATGA ATTAAAACT CAAGATTTTC AAAAAGTTCT	8820
ATTTTTTTCA AAGTAATAAA CTGACGGTAT AATAGGTAAA AAAAATTATT CTTTCTTGAT	8880
TTGTCAAAAT AAACAAATAA TTAGGAATAA TTAAAAAAT GGATAAATAA TTAACACGG	8940
AGGGAGCAAT ATGTTATCTT TAGCCTAATA ATATCTGATT AATGGCCACC CTAATTGATT	9000

GGATAGGAGA GGATAGACTT GCTTCCAAGT AACCCAAAAT ATAAAAAGTT GACAAAAGGG	9060
TGCTAAATTC GAGACACATG TAGTACTTAT ATAATTCATG TGC GGACTCG TTCTTTTGTA	9120
GTACTCCCTC CGTTCTATTT TATACGTCAC ATTTTACTT TATACTTTTA TTAAGAAATG	9180
ATGTAGTTTT ATCTTTCTAT TCTTATTTAA TGTTTTCTTA AGTCAATTTT ATAATAAATA	9240
ATGAATATAT TTTCAAGATT AATTAAC TACTATCAAGG GTATAATAGG TAAAATATGA	9300
TAATTTATAC ATAAATTTTA TAAAATGACA AGTATTGTGG TCCAAC TATT TATAGAAAGA	9360
AATGATATAT AAAATGGGAC GGAGGGCGTT ATAAAGTTGA CTTAAGAAAA CATTAATAA	9420
GGGTAGAAGG GTAAAATTAC ATTATTTCTT AATGTAAATG TAAAGTAAAA AGGTAACATA	9480
TAAAATGGAA AGGAGGGAGT AGTATTTTCT TGTTTTATTT TACGTGGCAC TCTATTCTCA	9540
TAATCCGTCT TTAAAAATGT CATTTTATTG TAATTGAAAA TAATTTAACT TAAAATTCTC	9600
CATCTACCCT TAATTAATGA AATGATTTAC AATTATATAA ATATATAAAA ATTGTTTTAG	9660
CCTATAATTT TCTAAAATCT TTTTTTTTCT CTTATACATC GTATTAAGTC AAACATAAAT	9720
GGAATGGACG GAGTATTTCT TTTATTTTTT TGTCACACCG CCCATATGTT TTCTCCCATC	9780
CCCCAGACCC CCACTATGTA TATTCACTCC TTAGTTGGAT CTGAATTTAG AGTTTAGAAG	9840
CTTCTATAAT AATTTTAGAT TAATATATAA TAATAATAAT AATAATTGAA CTTACAGTAT	9900
TAAATTTATG TGAATCTATA TATATTGTAT TGTAATTTTT TTAATTATAA TTTTAACCAA	9960
ATCAATAAAG CTATTCAGAT GTAAAAGTAT ATATTATGAT TTAACAACAA ATTTCTATAC	10020
GTCTTCCTAA GTTTTGATGC ATAATTTTCT AAAACTCATA AATTTCCAAG TGACTACTTC	10080
CAGTATTACA ATGAGAACTT ATGTTTCGTT ATGGATTTTC TTAGTGAATT AGTTTAATAA	10140
AATCAAAATG AAAAAAATC ATGTTTTATA ACATAAAATT TTCATTGATT CATGCGAAAA	10200
AAAAACATCT AGTTCTTATA GTGTGAAAAC TATTGAACTT ATGGGATGTA GCTGTATGGA	10260
AGTTCATCAA GTGGTAGCTC CTTGTACGCA ACTAGTGCTA CTTTTTATTG ACTAAAAGTT	10320
ATTTTCTAG	10329

## (2) INFORMATION FOR SEQ ID NO: 4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA oligonucleotide RG3

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Gpa2* encoding sequence of *S. tuberosum*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GGAGGCATCG GGAAAACAAC

20

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA oligonucleotide RG4

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Gpa2* encoding sequence of *S. tuberosum*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

TGCTAGAGGT AAYCCTCC

18

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA oligonucleotide RG5

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Gpa2* encoding sequence of *S. tuberosum*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

GATATGGTTG ACTCGGAATC AAG

23

## (2) INFORMATION FOR SEQ ID NO: 7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 bases
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA oligonucleotide RG6

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Gpa2* encoding sequence of *S. tuberosum*

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GAGTATGGAC CTCGATAGAG C

21

## REFERENCES

- Aarts, M.G.M., te Lintel Hekkert, B., Holub, E.B., Beynon, J.L., Stiekema, W.J. and Pereira, A. (1998). Identification of R-gene homologous DNA fragments genetically linked to disease loci in *Arabidopsis thaliana*. *Mol Plant-Microbe Interact.* 11: 251-258.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. (1990). Basic local alignment search tool. *J. Mol. Biol.* 215: 403-410.
- Arntzen, F.K., Visser, J.H.M. and Hoogendoorn, J. (1994). Inheritance, level and origin of resistance to *Globodera pallida* in the potato cultivar 'Multa', derived from *S. tuberosum* spp. *andigena* CPC1673. *Fundam. Appl. Nemat.* 16: 155-162.
- Baker, B., Zambryski, P., Staskawicz, B., and Dinesh-Kumar, S. P. (1997). Signaling in plant-microbe interactions. *Science* 276: 726-733.
- Bakker J., Bouwman-Smits L., and Gommers, F. J. (1992). Genetic relationships between *Globodera pallida* pathotypes in Europe assessed by using two dimensional gel electrophoresis of proteins. *Fundam Appl Nematol* 15: 481-490.
- Bakker, J., Folkertsma, R.T., Rouppe van der Voort, J.N.A.M., de Boer, J.M. and Gommers, F. (1993). Changing concepts and molecular approaches in the management of virulence genes in potato cyst nematodes. *Annu. Rev. Phytopathol.* 31: 169-190.
- Ballvora, A., Hesselbach, J., Niewohner, J., leister, D., Salamini, F and gebhardt, C. (1995). Marker enrichment and high-resolution map of the segment of potato chromosome VII harbouring the nematode resistance gene *Grol*. *Mol. Gen. Genet.* 249: 82-90.
- Barone, A., Ritter, E., Schachtschabel, U., Debener, T, Salamini, F. and Gebhardt, C. (1990). Localization by restriction length polymorphism mapping in potato of a major dominant gene conferring resistance to the potato cyst nematode *Globodera rostochiensis*. *Mol Gen. genet.* 224: 177-182.
- Bendahmane, A., Kanyuka, K., and Baulcombe, D. C. (1997). High-resolution genetical and physical mapping of the *Rx* gene for extreme resistance to potato virus X in tetraploid potato. *Theor. Appl. Genet.* 95: 153-162.
- Bent, A. F. Kunkel, B. N., Dahlbeck, D., Brown, K. L., Schmidt, R., Giraudat, J., Leung, J., and Staskawicz, B. J. (1994). *RPS2* of *Arabidopsis thaliana*: A leucine-rich repeat class of plant disease resistance genes. *Science* 265: 1856-1860.

- Burge, C. and Karlin, S. (1997).** Prediction of complete gene structure in human genomic DNA. *J. Mol. Biol.* **268**: 78-94.
- Büschges, R., Hollricher, K., Panstruga, R., Simons, G., Wolter, M., Frijters, A., van Daelen, R., van der Lee, T., Groenendijk, J., Topsch, S., Vos, P., Salamini, F., and Schultze-Lefert, P. (1997).** The barley *Mlo* gene: a novel control element of plant pathogen resistance. *Cell* **88**: 695-705.
- Chapman, S., Kavanagh, T. and Baulcombe, D. (1992).** Potato virus X as a vector for gene expression in plants. *Plant J.* **2**: 549-557.
- Chu, G. (1989).** Pulsed field electrophoresis in contour-clamped homogeneous electric fields for the resolution of DNA by size or topology. *Electrophoresis* **10**: 290-295.
- Crute, I. R., and Pink, D. A. C. (1996).** Genetics and utilization of pathogen resistance in plants. *Plant Cell* **8**:1747-1755.
- Dear, S. and Staden, R. (1991).** A sequence assembly and editing program for efficient management of large projects. *Nucleic Acids Res.* **14**: 3907-3911.
- De Jong, W., Forsyth, A., Leister, D., Gebhardt, C., and Baulcombe, D. C. (1997).** A potato hypersensitive resistance gene against potato virus X maps to a resistance gene cluster on chromosome 5. *Theor. Appl. Genet.* **95**: 246-252.
- Engbrecht, J., Brent, R. and Kaderbhai, M.A. (1997).** Large-scale preparation of plasmid DNA. In: Current Protocols in Molecular Biology, Vol. 1 (Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A. and Struhl, K., eds). John Wiley & Sons, Inc. pp. 1.6.1-1.6.2.
- Fenwick, D.W. (1940).** Methods for recovery and counting of cysts of *Heterodera schachtii* from soil. *J Helminth* **18**:155-172
- Folkertsma, R.T. (1997).** genetic diversity of the potato cyst nematode in the Netherlands. PhD thesis. Agricultural Univ. Wageningen, The Netherlands.
- Gebhardt, C., Mugniery, D., Ritter, E., Salamini, F. and Bonnel. (1993).** Identification of RFLP markers closely linked to the *Hl* gene conferring resistance to *Globodera rostochiensis* in potato. *Theor. Appl. Genet.* **85**: 541-544.
- Goulden, M.G., Köhm, B.A., Santa Cruz, S., kavanagh, T.A. and baulcombe, D. (1993).** A feature of the coat protein of potao virus X affects both induced virus resistance in potato and viral fitness. *Virology* **197**: 293-302.
- Heilig, J.S., Lech, K and Brent, R. (1997).** Large-scale preparation of plasmid DNA. In: Current Protocols in Molecular Biology, Vol. 1 (Ausubel, F.M., Brent, R., Kingston,

- R.E., Moore, D.D., Seidman, J.G., Smith, J.A. and Struhl, K., eds). John Wiley & Sons, Inc. pp. 1.7.1-1.7.3.
- Heungens, K., Mugniery, D., van Montagu, M., Gheysen, G. and Niebel, A. (1996).** A method to obtain disinfected *Globodera* infective juveniles directly from cysts. *Fundam. appl. Nemat.*, **19**, 91-93.
- Johal, G.S. and Briggs, S.P. (1992).** Reductase activity encoded by the Hm1 disease resistance gene in maize. *Science* **258**: 985-987.
- Konieczny, A. and Ausubel, F.M. (1993).** A procedure for mapping *Arabidopsis* mutations using co-dominant ecotype-specific PCR-based markers. *Plant J.* **4**: 403-410.
- 10 **Kreike, C. M., De Koning, J. R. A., Vinke, J. H., Van Ooijen, J. W., and Stiekema, W. J. (1994).** Quantitatively inherited resistance to *Globodera pallida* is dominated by one major locus in *Solanum spegazzinii*. *Theor. Appl. Genet.* **88**: 764-769.
- Lawrence, G. J., Finnegan, E. J., Ayliffe, M. A., and Ellis, J. G. (1995).** The *L6* gene for flax rust resistance is related to the *Arabidopsis* bacterial resistance gene *RPS2* and the tobacco viral resistance gene *N*. *Plant Cell* **7**:1195-1206.
- 15 **Lazo, G.R., Stein, P.A., and Ludwig, R.A. (1991).** A DNA transformation-competent *Arabidopsis* genomic library in *Agrobacterium*. *Bio/Technology* **9**: 963-967.
- Leister, D., Ballvora, A., Salamini F., and Gebhardt, C. (1996).** A PCR-based approach for isolating pathogen resistance genes from potato with potential for wide application in plants. *Nature Genetics* **14**: 421-429.
- 20 **Leonards-Schippers, C., Gieffers, W., Salamini, F., and Gebhardt, C. (1992).** The *R1* gene conferring race-specific resistance to *Phytophthora infestans* in potato is located on potato chromosome V. *Mol. Gen. Genet.* **233**: 378-383.
- Liu, Y-G and Whittier. (1994).** Rapid preparation of megabase plant DNA from nuclei in agarose plugs and microbeads. *Nucl. Acids Res.* **22**: 2168-2169.
- 25 **Manly, K. F. (1994).** Establishing genetic linkage using recombinant inbred lines with an abnormal segregation ratio. *Genetics* **136**:1434.
- Maruzyk, R. and Sergeant, A. (1980).** A simple method for dialysis of small volume samples. *Anal. Biochem.* **105**: 403-404.
- 30 **Mindrinos, M., Katagiri, F., Yu, G. L., and Ausubel, F. M. (1994).** The *A. thaliana* disease resistance gene *RPS2* encodes a protein a nucleotide-binding site and leucine - rich repeats. *Cell* **78**, 1089-1099.



- Ochman, H., Mehdora, M.M., Garza, D. and Hartl, D.L. (1990). Amplification of flanking sequences by inverse PCR. In *PCR Protocols* (Innis, M.A., Gelfand, D.H., Sninsky, J.J. and White, T.J., eds). San Diego: Academic Press, pp.219-227.
- Pineda, O., Bonierbale, M.W. and Plaisted, R.L. (1993). Identification of RFLP markers linked to the *Hl* gene conferring resistance to the potato cyst nematode *Globodera rostochiensis*. *Genome* 36: 153-156.
- Roupe van der Voort, J., Wolters, P., Folkertsma, R., Hutten, R., van Zandvoort, P., Vinke, H., Kanyuka, K., Bendahmane, A., Jacobsen, E., Janssen, R., and Bakker, J. (1997a). Mapping of the cyst nematode resistance locus *Gpa2* in potato using a strategy based on comigrating AFLP markers. *Theor. Appl. Genet.* 95: 874-880.
- Roupe van der Voort J.N.A.M., Van Zandvoort P., Eck H.J. van, Folkertsma, F.T., Hutten, R.C.B., Draaistra J., Gommers F.J., Jacobsen E., Helder J. and Bakker J. (1997b). Allele specificity of comigrating AFLP markers used to align genetic maps from different potato genotypes. *Mol Gen Genet* 255: 438-447.
- Roupe van der Voort, J., Lindeman, W., Folkertsma, R., Hutten, R., Overmars, H., van der Vossen, E., Jacobsen, E., and Bakker, J. (1998). A QTL for broad-spectrum resistance to cyst nematode species (genus *Globodera*) maps to a resistance gene cluster in potato. *Theor. Appl. Genet.* 96: 654-661.
- Sambrook J., Fritsch E.F. and Maniatis T. (1989). Molecular cloning: A Laboratory Manual (second edition). Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Säll, T. and Nilsson, N. O. (1994). The robustness of recombination frequency estimates in intercrosses with dominant markers. *Genetics* 137:589-596.
- Stam, P. (1993). Construction of integrated genetic linkage maps by means of a new computer package JoinMap. *P. Journal* 3: 739-744.
- Staskawicz, B. J., Ausubel, F. M., Baker, B. J., Ellis, J. G. and Jones, J. D. G. (1995). Molecular genetics of plant-disease resistance. *Science* 268:661-667.
- Suiter, K.A., Wendel, J. F., and Chase, J. S. (1983). LINKAGE-1: a Pascal computer program for the detection and analysis of genetic linkage. *J. Hered.* 74:203-204.
- Tanksley, S.D., Ganai, M.W., Prince, J.P., de Vicente, M.C., Bonierbale, M.W., Broun, P., Fulton, T.M., Giovannoni, J.J., Grandillo, S., Martin, G.B., Messeguer, R., Miller, J.C., Miller, L., Paterson, A.H., Pineda, O., Röder, M.S., Wing, R.A.,

**Wu, W. and Young, N.D. (1992).** High density molecular linkage maps of the tomato and potato genomes. *Genetics* **132**: 1141-1160.

**Van Engelen, F.A., Molthoff, J.W., Conner, A.J, Nap, J-P., Pereira, A. and Stiekema, W.J. (1995).** pBIBPLUS: an improved plant transformation vector based on pBIN19. *Transgenic Research* **4**: 288-290.

**Visser, R.G.F. (1991).** Regeneration and transformation of potato by *Agrobacterium tumefaciens*. In: Plant tissue culture manual. Kluwer Academic Publishers, Dordrecht Boston London, B5:1-9.

**Whitham, S. Dinesh-Kumar, S.P., Choi, D., Hehl, R., Corr, C. and Baker, B. (1994).** The product of the tobacco mosaic virus resistance gene N - similarity to Toll and the interleukin-1 receptor. *Cell* **78**: 1101-1115.

**Young, N., and Tanksley, S. D. (1992).** Restriction fragment length polymorphism maps and the concept of graphical genotypes. *Theor. Appl. Genet.* **77**:95-101.

## CLAIMS

1. A recombinant nucleic acid sequence providing resistance to infection by a  
5 phytopathogenic nematode of the genus *Globodera* when introduced into a host plant, said  
host plant prior to introduction being susceptible to infection with the phytopathogenic  
nematode, said introduction occurring in such a way that said nucleic acid sequence is  
expressed in the host plant, the nucleic acid sequence being that of SEQ ID NO.1.
- 10 2. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence  
according to claim 1, said homologue also providing the resistance, said homologue being  
a nucleic acid sequence encoding the amino acid sequence of SEQ ID NO.1.
- 15 3. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence  
according to claim 1, said homologue also providing the resistance, said homologue being  
a nucleic acid sequence exhibiting more than 70% homology at nucleic acid level with  
SEQ ID NO. 1.
- 20 4. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence  
according to claim 1, said homologue also providing the resistance, said homologue being  
a nucleic acid sequence exhibiting more than 75% homology at nucleic acid level with  
SEQ ID NO. 1.
- 25 5. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence  
according to claim 1, said homologue also providing the resistance, said homologue being  
a nucleic acid sequence exhibiting more than 80% homology at nucleic acid level with  
SEQ ID NO. 1.
- 30 6. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence  
according to claim 1, said homologue also providing the resistance, said homologue being  
a nucleic acid sequence exhibiting more than 85% homology at nucleic acid level with  
SEQ ID NO. 1.

7. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence according to claim 1, said homologue also providing the resistance, said homologue being a nucleic acid sequence exhibiting more than 90% homology at nucleic acid level with SEQ ID NO. 1.

5

8. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence according to claim 1, said homologue also providing the resistance, said homologue being a nucleic acid sequence exhibiting more than 95% homology at nucleic acid level with SEQ ID NO. 1.

10

9. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence according to claim 1 or being a homologue according to any of the claims 2-8, said homologue also providing the resistance, said homologue being a nucleic acid sequence capable of hybridising under normal to stringent conditions to the nucleic acid sequence of SEQ ID NO. 1.

15

10. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence according to claim 1 or being a homologue according to any of claims 2-9, said homologue also providing the resistance, said homologue being a nucleic acid sequence encoding a deletion, insertion or substitution mutant of the amino acid sequence of SEQ ID NO. 1.

20

11. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence according to claim 1 or being a homologue according to any of claims 2-10, said homologue also providing the resistance, said homologue being a nucleic acid sequence encoding a deletion, insertion or substitution variant as occurs in nature of the amino acid sequence of SEQ ID NO. 1.

25

12. A recombinant nucleic acid sequence according to any of the preceeding claims, said nucleic acid sequence further comprising at least one intron.

30

13. A recombinant nucleic acid sequence according to claim 12 comprising at least one intron of SEQ ID NO. 2.

14. A recombinant nucleic acid according to any of the preceeding claims being the genomic insert of pBINRGH2.
- 5 15. A recombinant nucleic acid sequence according to any of the preceeding claims, said nucleic acid sequence being identical to that present in the genetic material of a species of the family Solanaceae, preferably a species of the genus *Solanum*.
- 10 16. A recombinant nucleic acid sequence according to any of the preceeding claims, said nucleic acid sequence being identical to that present in the genetic material of a potato, preferably on chromosome 4, 5, 6, 7, 9, 11 or 12.
- 15 17. A recombinant nucleic acid sequence according to any of the preceeding claims, said nucleic acid sequence being identical to that present in the genetic material of potato locus *Gpa2*.
- 20 18. A recombinant nucleic acid sequence being a homologue of the nucleic acid sequence according to claim 1, said homologue also providing the resistance, said homologue being a fragment of the nucleic acid sequence according to any of claims 14-17.
- 25 19. A genetic construct comprising a nucleic acid sequence according to any of the preceeding claims said sequence being operably linked to a regulatory region for expression.
- 20 20. A genetic construct according to claim 19 wherein the regulatory region is a *Gpa2* regulatory region.
- 30 21. A genetic construct according to any of claims 19 or 20 wherein the regulatory region corresponds to that present in the sequence of nucleotides 1-4874 of SEQ ID NO. 3.
22. A genetic construct according to any of claims 19-21, wherein the regulatory region corresponds to that of nucleotides 1-4874 of SEQ ID NO.3.

23. A genetic construct according to any of the preceeding claims 19-22, wherein the regulatory region comprises a promoter functionally connected to a nucleic acid sequence as defined in any of the claims 1-18, said promoter being able to control the transcription of said nucleic acid sequence in a plant cell.

5

24. A vector which carries a nucleic acid according to any of the claims 1-18, or a genetic construct according to any of the claims 19-23.

25. A vector according to claim 24 capable of replicating in a host organism.

10

26. A vector capable of expressing the nucleic acid according to any of the claims 1-19, or a genetic construct according to any of the claims 19-23.

15

27. A vector according to any of claims 24-26 constructed to function in a host organism selected from the group consisting of a micro-organism, plant cell, plant, seed, seedling, plant part and protoplast.

20

28. A vector according to any of claims 24-27 constructed to function in a host organism selected from the group consisting of a micro-organism, plant cell, seed, seedling, plant part and protoplast.

25

29. A vector according to any of claims 24-28 constructed to function in a host organism selected from the group consisting of a micro-organism, plant cell, plant part and protoplast.

30. A vector according to any of claims 24-29 constructed to function in a host organism selected from the group consisting of a plant, plant cell, plant part, seed, seedling and protoplast.

30

31. A host organism selected from the group consisting of a micro-organism, plant cell, plant, seed, seedling, plant part and protoplast, harbouring a vector as defined in any of claims 24-30 and/or a genetic construct according to any of the claims 19-23.

32. A host organism selected from the group consisting of a micro-organism, plant cell, seed, seedling, plant part and protoplast, harbouring a vector as defined in any of claims 24-30 and/or a genetic construct according to any of the claims 19-23.

5 33. A host organism selected from the group consisting of a micro-organism, plant cell, plant part and protoplast, harbouring a vector as defined in any of claims 24-30 and/or a genetic construct according to any of the claims 19-23.

10 34. A host organism selected from the group consisting of a plant cell, plant, seed, seedling, plant part and protoplast, harbouring a vector as defined in any of claims 24-30 and/or a genetic construct according to any of the claims 19-23.

15 35. A host organism according to any preceeding claim 31-34 which is capable of replicating or expressing the nucleic acid sequence or the genetic construct of the vector and/or a genetic construct according to any of the claims 19-23.

20 36. A process for producing a genetically transformed or transfected host organism having increased resistance to phytopathogenic nematodes of the genus *Globodera* as compared to the host organism prior to the transformation, said process comprising transferring a genetic construct according to any of the claims 19-23 and/or a vector according to any of claims 24-30 into the host organism so that it's genetic material comprises the genetic construct and subsequently regenerating the host organism into a genetically transformed plant part.

25 37. A process according to claim 36 for producing a genetically transformed plant having increased resistance to phytopathogenic nematodes of the genus *Globodera* as compared to a corresponding plant prior to the transformation, said process comprising transferring a genetic construct according to any of the claims 19-23 and/or a vector according to any of claims 24-30 into the host organism so that it's genetic material comprises the genetic  
30 construct and/or a vector according to any of claims 19-23 and subsequently regenerating the host organism into a genetically transformed plant, said host organism being selected from the group consisting of a plant cell, plant, seed, seedling, plant part and protoplast of the plant type to be rendered resistant.

38. A process according to claim 36 or 37 wherein said nematodes are selected from the group consisting of *Globodera pallida* and *Globodera rostochiensis*.

5 39. A process according to any of claims 36-38, wherein said host organism to be transformed is selected from a plant type of the family Solanaceae.

40. A process according to any of claims 36-39, wherein said host organism to be transformed is selected from a plant type of the genus *Solanum*.

10

41. A process according to any of claims 36-40, wherein said host organism to be transformed is selected from a plant type of the species *Solanum tuberosum*.

15

42. A process for isolating or detecting a nucleic acid sequence according to any of claims 1-18, comprising the screening of genomic nucleic acid of a plant with a nucleic acid sequence according to any of claims 1-18 or a fragment thereof as probe or primer, said probe or primer being at least 16 nucleotides in length, the identification of positive clones which hybridize to said probe and the isolation of said positive clones and the isolation of the nucleic acid sequence therefrom.

20

43. A process for isolating or detecting a nucleic acid sequence according to claims 1-18, comprising the screening of a genomic library of a plant with a nucleic sequence according to seq id no 3 or a fragment thereof as probe, said probe or primer being at least 16 nucleotides in length, the identification of positive clones which hybridize to said probe or primer and the isolation of said positive clones and the isolation of the nucleic acid sequence therefrom.

25

44. A process for isolating or detecting a nucleic acid sequence according to claims 1-18, comprising the screening of a cDNA library of a plant with the encoding portion of a nucleic acid sequence according to any of claims 1-18 or a fragment thereof as probe or primer, said probe or primer being at least 16 nucleotides in length, the identification of positive clones which hybridize to said probe or primer and the isolation of said positive clones and the isolation of the nucleic acid sequence therefrom.

30



45. A process for isolating or detecting a nucleic acid sequence according to claims 1-18, comprising the screening of a cDNA library of a plant with the encoding portion of a nucleic acid sequence according to SEQ ID NO. 1 or a fragment thereof as probe or  
5 primer, said probe or primer being at least 16 nucleotides in length, the identification of positive clones which hybridize to said probe and the isolation of said positive clones and the isolation of the nucleic acid sequence therefrom.

46. A process according to any of claims 42-45, wherein the probe is comprised within the  
10 sequence of SEQ ID NO.1, SEQ ID NO.2 or SEQ ID NO.3.

47. A process for isolating or detecting a nucleic acid sequence according to any of claims 1-18, using a nucleic acid amplification reaction such as the Polymerase Chain Reaction and at least one primer corresponding to or being complementary to the nucleic acid  
15 sequence according to any of claims 1-18 or a fragment thereof as primer, said primer being at least 16 nucleotides in length.

48. A process for isolating or detecting a nucleic acid sequence according to any of claims 1-18, using a nucleic acid amplification reaction such as the Polymerase Chain Reaction and at least one primer corresponding to or being complementary to the nucleic acid  
20 sequence of of SEQ ID NO.1, SEQ ID NO.2 or SEQ ID NO.3 or a fragment thereof as primer, said primer being at least 16 nucleotides in length.

49. A process according to any of claims 42-48 wherein said probe or primer comprises a  
25 nucleic acid sequence encoding at least part of the amino acid sequence of the NBS sequence of *Gpa2*.

50. A process according to any of claims 42-49, wherein said probe or primer comprises a nucleic acid sequence encoding at least part of the amino acid sequence of the NBS  
30 sequence of *Gpa2*, at least part having the following sequence GGIGKTT or GGLPLA.

51. A process according to any of claims 42-50, wherein said probe or primer comprises parts of the NBS sequence of *Gpa2* and at least one codon of a 5' and/or 3' overhanging

portion corresponding to the respective 5' and/or 3' adjacent amino acids of the specified NBS sequence of *Gpa2*.

52. A process according to any of claims 42-51, wherein said probe or primer comprises  
5 parts of the NBS sequence of *Gpa2* and at least one codon of a 5' and/or 3' overhanging  
portion corresponding to the respective 5' and/or 3' adjacent amino acids of the specified  
NBS sequence of *Gpa2* of SEQ ID NO.1.

53. A process according to any of claims 42-52, wherein said probe or primer corresponds  
10 to a sequence selected from SEQ ID NO.4, SEQ ID NO.5, SEQ ID NO.6 and/or SEQ ID  
NO.7.

54. A polypeptide having an amino acid sequence provided in SEQ ID NO.1 or being a  
homologue of said amino acid sequence, said homologue being a substitution, insertion or  
15 deletion mutant possessing nematode resistance against a nematode of the genus  
*Globodera*.

55. A polypeptide encoded by a sequence according to any of the claims 1-18.

20 56. A process for producing a polypeptide having an amino acid sequence provided in  
SEQ ID NO.1 or a homologue of said amino acid sequence, said homologue being a  
substitution, insertion or deletion mutant possessing nematode resistance against a  
nematode of the genus *Globodera*, said process comprising expressing a recombinant  
nucleic acid sequence according to any of the claims 1-18 or genetic construct according  
25 to any of claims 19-23 and optionally isolating said polypeptide, said expression occurring  
in a host organism according to any of claims 31-35.

57. A process for producing a polypeptide having an amino acid sequence provided in  
SEQ ID NO.1 or a homologue of said amino acid sequence, said homologue being a  
30 substitution, insertion or deletion mutant possessing nematode resistance against a  
nematode of the genus *Globodera*, said process comprising the expression of a  
recombinant nucleic acid sequence according to any of the claims 1-18 or genetic

construct according to any of claims 19-23 and isolating said polypeptide, said expression occurring in a host organism according to any of claims 31-35.

58. A nematocide composition comprising as active ingredient a polypeptide according to claim 54 or 55 or produced according to claim 56 or 57 or a host organism expressing such a polypeptide, such a host organism being defined in any of claims 31-35 in a formulation suitable for application as nematocide to a plant and optionally comprising other ingredients required for rendering the polypeptide suitable for application as a nematocide.

10

59. A nematocide composition according to claim 58 comprising the polypeptide in a slow release dosage form.

15

60. A nematocide composition according to 58 or 59 comprising instructions for application as nematocide.

20

61. A nucleic acid sequence comprising at least 16 contiguous nucleotides corresponding to or complementary to the *Gpa2* sequence, with the *proviso* that when such an oligonucleotide comprises part or all of the NBS encoding sequence, the oligonucleotide also comprises at least one codon of a 5' and/or 3' overhanging portion corresponding to the respective 5' and/or 3' adjacent amino acids of the specified NBS sequence of *Gpa2*.

25

62. A nucleic acid sequence according to claim 61, wherein the *Gpa2* sequence is comprised within the sequence of SEQ ID NO.1, 2 or 3.

30

63. A nucleic acid sequence according to claim 61 or 62, wherein sequence length is at least 50 nucleotides, suitably more than 100 nucleotides and is suitable for use as probe or primer in a nucleic acid assay.

64. A nucleic acid sequence according to any of claims 61-63, being selected from any of the sequences SEQ ID NOs. 4, 5, 6 and/or 7.

65. A combination of at least 2 primers according to any of claims 61-64.

66. Antibody raised against a polypeptide of claim 55 or a polypeptide produced by a process according to claim 56 or 57.

- 5 67. A diagnostic kit for assessing the presence of nematode resistance of a plant to infection by a phytopathogenic nematode of the genus *Globodera*, said kit comprising at least one nucleic acid sequence according to any of claims 61-64 as a probe or primer for screening of nucleic acid from a plant or plant part to be tested and/or a combination of primers according to claim 65 and/or an antibody according to claim 66.

10


68. A process for diagnosing whether a plant is resistant to a phytopathogenic nematode of the genus *Globodera*, said process comprising detecting the presence of a nucleic acid sequence according to any of claims 1-18, genetic construct according to any of claims 19-23, vector according to any of 24-30 or a polypeptide according to claim 55 in plant material of a plant to be tested.
- 15

69. A process for diagnosing whether a plant is resistant to a phytopathogenic nematode of the genus *Globodera*, said process comprising carrying out a process according to any of claims 42-53 and/or applying a diagnostic kit according to claim 67.

20


70. A process for protecting plants said process comprising the introduction of the nucleic acid sequence according to any of claims 1-18, the genetic construct according to any of claims 19-23, the vector according to any of 24-30 in plant material of a plant to be protected.

1947



2/16

fig-2



3/16

Fig. 3a (1)

atg gct tat gct gct gtt act tcc ctt atg aga acc ata cat caa tca	48
Met Ala Tyr Ala Ala Val Thr Ser Leu Met Arg Thr Ile His Gln Ser	
1 5 10 15	
atg gaa ctt act gga tgt gat ttg caa ccg ttt tat gaa aag ctc aaa	96
Met Glu Leu Thr Gly Cys Asp Leu Gln Pro Phe Tyr Glu Lys Leu Lys	
20 25 30	
tct ttg aga gct att ctg gag aaa tcc tgc aat ata atg ggc gat cat	144
Ser Leu Arg Ala Ile Leu Glu Lys Ser Cys Asn Ile Met Gly Asp His	
35 40 45	
gag ggg tta aca atc ttg gaa gtt gaa atc ata gag gta gca tac aca	192
Glu Gly Leu Thr Ile Leu Glu Val Glu Ile Ile Glu Val Ala Tyr Thr	
50 55 60	
aca gaa gat atg gtt gac tcg gaa tca aga aat gtt ttt tta gca cgg	240
Thr Glu Asp Met Val Asp Ser Glu Ser Arg Asn Val Phe Leu Ala Arg	
65 70 75 80	
aat gtg ggg aaa aga agc agg gct atg tgg ggg att ttt ttc gtc ttg	288
Asn Val Gly Lys Arg Ser Arg Ala Met Trp Gly Ile Phe Phe Val Leu	
85 90 95	
gaa caa gca cta gaa tgc att gat tcc acc gtg aaa cag tgg atg gca	336
Glu Gln Ala Leu Glu Cys Ile Asp Ser Thr Val Lys Gln Trp Met Ala	
100 105 110	
aca tcg gac agc atg aaa gat cta aaa cca caa act agc tca ctt gtc	384
Thr Ser Asp Ser Met Lys Asp Leu Lys Pro Gln Thr Ser Ser Leu Val	
115 120 125	
agt tta cct gaa cat gat gtt gag cag ccc gag aat ata atg gtt ggc	432
Ser Leu Pro Glu His Asp Val Glu Gln Pro Glu Asn Ile Met Val Gly	
130 135 140	
cgt gaa aat gaa ttt gag atg atg ctg gat caa ctt gct aga gga gga	480
Arg Glu Asn Glu Phe Glu Met Met Leu Asp Gln Leu Ala Arg Gly Gly	
145 150 155 160	
agg gaa cta gaa gtt gtc tca atc gta ggg atg gga ggc atc ggg aaa	528
Arg Glu Leu Glu Val Val Ser Ile Val Gly Met Gly Gly Ile Gly Lys	
165 170 175	
aca act ttg gct gca aaa ctc tat agt gat cct tac att atg tct cga	576
Thr Thr Leu Ala Ala Lys Leu Tyr Ser Asp Pro Tyr Ile Met Ser Arg	
180 185 190	
ttt gat att cgt gca aaa gca act gtt tca caa gag tat tgt gtg aga	624
Phe Asp Ile Arg Ala Lys Ala Thr Val Ser Gln Glu Tyr Cys Val Arg	
195 200 205	

4/16

Fig. 3a (2)

aat gta ctc cta ggc ctt ctt tct ttg aca agt gat gaa cct gat tat	672
Asn Val Leu Leu Gly Leu Leu Ser Leu Thr Ser Asp Glu Pro Asp Tyr	
210 215 220	
cag cta gcg gac caa ctg caa aag cat ctg aaa ggc agg aga tac ttg	720
Gln Leu Ala Asp Gln Leu Gln Lys His Leu Lys Gly Arg Arg Tyr Leu	
225 230 235 240	
gta gtc att gat gac ata tgg act aca gaa gct tgg gat gat ata aaa	768
Val Val Ile Asp Asp Ile Trp Thr Thr Glu Ala Trp Asp Asp Ile Lys	
245 250 255	
cta tgt ttc cca gac tgc gat aat gga agc aga ata ctc ctg act act	816
Leu Cys Phe Pro Asp Cys Asp Asn Gly Ser Arg Ile Leu Leu Thr Thr	
260 265 270	
cgg aat gtg gaa gtg gct gaa tat gct agc tca ggt aag cct cct cat	864
Arg Asn Val Glu Val Ala Glu Tyr Ala Ser Ser Gly Lys Pro Pro His	
275 280 285	
cac atg cgc ctc atg aat ttt gac gaa agt tgg aat tta cta cac aaa	912
His Met Arg Leu Met Asn Phe Asp Glu Ser Trp Asn Leu Leu His Lys	
290 295 300	
aag atc ttt gaa aaa gaa ggt tct tat tct cct gaa ttt gaa aat att	960
Lys Ile Phe Glu Lys Glu Gly Ser Tyr Ser Pro Glu Phe Glu Asn Ile	
305 310 315 320	
ggg aaa caa att gca tta aaa tgt gga ggg tta cct cta gca att act	1008
Gly Lys Gln Ile Ala Leu Lys Cys Gly Gly Leu Pro Leu Ala Ile Thr	
325 330 335	
ttg att gct gga ctt ctc tcc aaa atc agt aaa aca ttg gat gag tgg	1056
Leu Ile Ala Gly Leu Leu Ser Lys Ile Ser Lys Thr Leu Asp Glu Trp	
340 345 350	
caa aat gtt gcg gag aat gta cgt tgc gtg gta agc aca gat ctt gaa	1104
Gln Asn Val Ala Glu Asn Val Arg Ser Val Val Ser Thr Asp Leu Glu	
355 360 365	
gca aaa tgc atg aga gtg ttg gct ttg agt tac cat cac ttg cct tct	1152
Ala Lys Cys Met Arg Val Leu Ala Leu Ser Tyr His His Leu Pro Ser	
370 375 380	
cac cta aaa ccg tgt ttt ctg tat ttt gca att ttc gca gag gat gaa	1200
His Leu Lys Pro Cys Phe Leu Tyr Phe Ala Ile Phe Ala Glu Asp Glu	
385 390 395 400	
cgg att tat gta aat aaa ctt gtt gag tta tgg gcc gta gag ggg ttt	1248
Arg Ile Tyr Val Asn Lys Leu Val Glu Leu Trp Ala Val Glu Gly Phe	
405 410 415	



5/16

Fig. 3a (3)

ttg aat gaa gaa gag gga aaa agc ata gaa gag gtg gca gaa aca tgt	1296
Leu Asn Glu Glu Glu Gly Lys Ser Ile Glu Glu Val Ala Glu Thr Cys	
420 425 430	
ata aac gaa ctt gta gat aga agt cta att tct atc cac aat gtg agt	1344
Ile Asn Glu Leu Val Asp Arg Ser Leu Ile Ser Ile His Asn Val Ser	
435 440 445	
ttt gat ggg gaa aca cag aga tgt gga atg cat gat gtg acc cgt gaa	1392
Phe Asp Gly Glu Thr Gln Arg Cys Gly Met His Asp Val Thr Arg Glu	
450 455 460	
ctc tgt ttg agg gaa gct cga aac atg aat ttt gtg aat gtt atc aga	1440
Leu Cys Leu Arg Glu Ala Arg Asn Met Asn Phe Val Asn Val Ile Arg	
465 470 475 480	
gga aag agt gat caa aat tca tgt gca caa tcc atg cag tgt tcc ttt	1488
Gly Lys Ser Asp Gln Asn Ser Cys Ala Gln Ser Met Gln Cys Ser Phe	
485 490 495	
aag agt cga agt cgg atc agt atc cat aat gag gaa gaa ttg gtt tgg	1536
Lys Ser Arg Ser Arg Ile Ser Ile His Asn Glu Glu Glu Leu Val Trp	
500 505 510	
tgt cgt aac agc gag gct cat tct atc atc acg ttg tgt ata ttc aaa	1584
Cys Arg Asn Ser Glu Ala His Ser Ile Ile Thr Leu Cys Ile Phe Lys	
515 520 525	
tgc gtc aca ctg gaa ttg tct ttc aag cta gta aga gta cta gat ctt	1632
Cys Val Thr Leu Glu Leu Ser Phe Lys Leu Val Arg Val Leu Asp Leu	
530 535 540	
ggt ttg act aca tgc cca att ttt ccc agt gga gta ctt tct cta att	1680
Gly Leu Thr Thr Cys Pro Ile Phe Pro Ser Gly Val Leu Ser Leu Ile	
545 550 555 560	
cat ttg aga tac cta tct ttg cgt ttt aat cct cgc tta cag cag tat	1728
His Leu Arg Tyr Leu Ser Leu Arg Phe Asn Pro Arg Leu Gln Gln Tyr	
565 570 575	
cga gga tcg aaa gaa gct gtt ccc tca tca ata ata gac att cct cta	1776
Arg Gly Ser Lys Glu Ala Val Pro Ser Ser Ile Ile Asp Ile Pro Leu	
580 585 590	
tcg ata tca agc cta tgc tat ctg caa act ttt aaa ctt tac cat cca	1824
Ser Ile Ser Ser Leu Cys Tyr Leu Gln Thr Phe Lys Leu Tyr His Pro	
595 600 605	
ttt ccc aat tgt tat cct ttc ata tta cca tcg gaa att ttg aca atg	1872
Phe Pro Asn Cys Tyr Pro Phe Ile Leu Pro Ser Glu Ile Leu Thr Met	
610 615 620	

6/16

Fig. 3a (4)

cca	caa	ttg	agg	aag	ctg	tgt	atg	ggc	tgg	aat	tac	ttg	cgg	agt	cat	1920
Pro	Gln	Leu	Arg	Lys	Leu	Cys	Met	Gly	Trp	Asn	Tyr	Leu	Arg	Ser	His	
625					630					635					640	
gag	cct	aca	gag	aac	aga	ttg	gtt	ttg	aaa	agt	ttg	caa	tgc	ctc	aat	1968
Glu	Pro	Thr	Glu	Asn	Arg	Leu	Val	Leu	Lys	Ser	Leu	Gln	Cys	Leu	Asn	
				645					650					655		
gaa	ttg	aat	cct	cgg	tat	tgt	aca	ggg	tct	ttt	tta	aga	cta	ttt	ccc	2016
Glu	Leu	Asn	Pro	Arg	Tyr	Cys	Thr	Gly	Ser	Phe	Leu	Arg	Leu	Phe	Pro	
			660					665					670			
aat	tta	aag	aag	ttg	gaa	gta	ttt	ggc	gtc	aaa	gag	gac	ttt	cgc	aat	2064
Asn	Leu	Lys	Lys	Leu	Glu	Val	Phe	Gly	Val	Lys	Glu	Asp	Phe	Arg	Asn	
		675					680					685				
cac	aag	gac	ctg	tat	gat	ttt	cgc	tac	tta	tat	cag	ctc	gag	aaa	ttg	2112
His	Lys	Asp	Leu	Tyr	Asp	Phe	Arg	Tyr	Leu	Tyr	Gln	Leu	Glu	Lys	Leu	
	690					695					700					
gca	ttt	agt	act	tat	tat	tca	tct	tct	gct	tgc	ttt	cta	aaa	aac	act	2160
Ala	Phe	Ser	Thr	Tyr	Tyr	Ser	Ser	Ser	Ala	Cys	Phe	Leu	Lys	Asn	Thr	
705					710					715					720	
gca	cct	tta	ggt	tct	act	ccg	caa	gat	cct	ctg	agg	ttt	cag	atg	gaa	2208
Ala	Pro	Leu	Gly	Ser	Thr	Pro	Gln	Asp	Pro	Leu	Arg	Phe	Gln	Met	Glu	
			725						730					735		
aca	ttg	cac	tta	gag	act	cat	tcc	agg	gca	act	gca	cct	cca	act	gat	2256
Thr	Leu	His	Leu	Glu	Thr	His	Ser	Arg	Ala	Thr	Ala	Pro	Pro	Thr	Asp	
			740					745					750			
gtt	cca	act	ttc	ctc	tta	cct	cct	ccg	gat	tgt	ttt	cca	caa	aac	ctt	2304
Val	Pro	Thr	Phe	Leu	Leu	Pro	Pro	Pro	Asp	Cys	Phe	Pro	Gln	Asn	Leu	
		755					760					765				
aag	agt	tta	act	ttt	agc	gga	gat	ttc	ttt	ttg	gca	tgg	aag	gat	ttg	2352
Lys	Ser	Leu	Thr	Phe	Ser	Gly	Asp	Phe	Phe	Leu	Ala	Trp	Lys	Asp	Leu	
	770					775					780					
agc	att	gtt	ggt	aaa	tta	ccc	aaa	ctc	gag	gtc	ctt	caa	cta	tca	cac	2400
Ser	Ile	Val	Gly	Lys	Leu	Pro	Lys	Leu	Glu	Val	Leu	Gln	Leu	Ser	His	
785					790					795					800	
aat	gcc	ttc	aaa	ggc	gag	gag	tgg	gaa	gta	gtt	gag	gaa	ggg	ttt	cct	2448
Asn	Ala	Phe	Lys	Gly	Glu	Glu	Trp	Glu	Val	Val	Glu	Glu	Gly	Phe	Pro	
				805					810					815		
cac	ttg	aag	ttc	ttg	ttt	ctg	gat	agc	ata	tac	att	cgg	tac	tgg	aga	2496
His	Leu	Lys	Phe	Leu	Phe	Leu	Asp	Ser	Ile	Tyr	Ile	Arg	Tyr	Trp	Arg	
			820					825					830			

7/16

**Fig. 3a (5)**

gct agt agt gat cac ttt cca tac ctt gaa cga ctt ttt ctt agc gat	2544
Ala Ser Ser Asp His Phe Pro Tyr Leu Glu Arg Leu Phe Leu Ser Asp	
835 840 845	
tgc ttt tat ttg gat tca atc cct cga gat ttt gca gat ata acc aca	2592
Cys Phe Tyr Leu Asp Ser Ile Pro Arg Asp Phe Ala Asp Ile Thr Thr	
850 855 860	
cta gct ctt att gat ata ttt cgc tgc caa caa tct gtt ggg aat tcc	2640
Leu Ala Leu Ile Asp Ile Phe Arg Cys Gln Gln Ser Val Gly Asn Ser	
865 870 875 880	
gcc aag caa att caa cag gac att caa gac aac tat gga agc tct atc	2688
Ala Lys Gln Ile Gln Gln Asp Ile Gln Asp Asn Tyr Gly Ser Ser Ile	
885 890 895	
gag gtc cat act cgt tat ctt tat cga aat gga gca ttt ttg gta gtg	2736
Glu Val His Thr Arg Tyr Leu Tyr Arg Asn Gly Ala Phe Leu Val Val	
900 905 910	
tga	2739
*	

8/16

**Fig. 3b (1)**

ATGGCTTATG CTGCTGTTAC TTCCCTTATG AGAACCATAC ATCAATCAAT GGAACCTACT	60
GGATGTGATT TGCAACCGTT TTATGAAAAG CTCAAATCTT TGAGAGCTAT TCTGGAGAAA	120
TCCTGCAATA TAATGGGCGA TCATGAGGGG TTAACAATCT TGGAAGTTGA AATCATAGAG	180
GTAGCATACA CAACAGAAGA TATGGTTGAC TCGGAATCAA GAAATGTTTT TTTAGCACGG	240
AATGTGGGGA AAAGAAGCAG GGCTATGTGG GGGATTTTTT TCGTCTTGGA ACAAGCACTA	300
GAATGCATTG ATTCCACCGT GAAACAGTGG ATGGCAACAT CGGACAGCAT GAAAGATCTA	360
AAACCACAAA CTAGCTCACT TGTCAGTTTA CCTGAACATG ATGTTGAGCA GCCCCAGAAT	420
ATAATGGTTG GCCGTGAAAA TGAATTTGAG ATGATGCTGG ATCAACTTGC TAGAGGAGGA	480
AGGGAAGTAG AAGTTGTCTC AATCGTAGGG ATGGGAGGCA TCGGGAAAAC AACTTTGGCT	540
GCAAACTCT ATAGTGATCC TTACATTATG TCTCGATTTG ATATTCGTGC AAAAGCAACT	600
GTTTCACAAG AGTATTGTGT GAGAAATGTA CTCCTAGGCC TTCTTTCTTT GACAAGTGAT	660
GAACCTGATT ATCAGCTAGC GGACCAACTG CAAAAGCATC TGAAAGGCAG GAGATACTTG	720
GTAGTCATTG ATGACATATG GACTACAGAA GCTTGGGATG ATATAAACT ATGTTTCCCA	780
GACTGCGATA ATGGAAGCAG AATACTCCTG ACTACTCGGA ATGTGGAAGT GGCTGAATAT	840
GCTAGCTCAG GTAAGCCTCC TCATCACATG CGCCTCATGA ATTTTGACGA AAGTTGGAAT	900
TTACTACACA AAAAGATCTT TGAAAAAGAA GGTTCCTTATT CTCCTGAATT TGAAAATATT	960
GGGAAACAAA TTGCATTAAA ATGTGGAGGG TTACCTCTAG CAATTACTTT GATTGCTGGA	1020
CTTCTCTCCA AAATCAGTAA AACATTGGAT GAGTGGCAAA ATGTTGCGGA GAATGTACGT	1080
TCGGTGGTAA GCACAGATCT TGAAGCAAAA TGCATGAGAG TGTGGCTTT GAGTTACCAT	1140
CACTTGCCTT CTCACCTAAA ACCGTGTTTT CTGTATTTTG CAATTTTCGC AGAGGATGAA	1200
CGGATTTATG TAAATAAACT TGTGAGTTA TGGGCCGTAG AGGGGTTTTT GAATGAAGAA	1260
GAGGGAAAAA GCATAGAAGA GGTGGCAGAA ACATGTATAA ACGAACTTGT AGATAGAAGT	1320
CTAATTTCTA TCCACAATGT GAGTTTTGAT GGGGAAACAC AGAGATGTGG AATGCATGAT	1380
GTGACCCGTG AACTCTGTTT GAGGGAAGCT CGAAACATGA ATTTTGTGAA TGTTATCAGA	1440
GGAAAGAGTG ATCAAAATTC ATGTGCACAA TCCATGCAGT GTTCCTTTAA GAGTCGAAGT	1500
CGGATCAGTA TCCATAATGA GGAAGAATTG GTTTGGTGTC GTAACAGCGA GGCTCATTCT	1560
ATCATCACGT TGTGTATATT CAAATGCGTC AACTGGAAT TGTCTTTCAA GCTAGTAAGA	1620

9/16

**Fig. 3b (2)**

GTACTAGATC TTGGTTTGAC TACATGCCCA ATTTTCCCA GTGGAGTACT TTCTCTAATT	1680
CATTTGAGAT ACCTATCTTT GCGTTTAAAT CCTCGCTTAC AGCAGTATCG AGGATCGAAA	1740
GAAGCTGTTT CCTCATCAAT AATAGACATT CCTCTATCGA TATCAAGCCT ATGCTATCTG	1800
CAAACCTTTTA AACTTTACCA TCCATTTCCC AATTGTTATC CTTTCATATT ACCATCGGAA	1860
ATTTTGACAA TGCCACAATT GAGGAAGCTG TGTATGGGCT GGAATTACTT GCGGAGTCAT	1920
GAGCCTACAG AGAACAGATT GGTTTTGAAA AGTTTGCAAT GCCTCAATGA ATTGAATCCT	1980
CGGTATTGTA CAGGGTCTTT TTTAAGACTA TTTCCCAATT TAAAGAAGTT GGAAGTATTT	2040
GGCGTCAAAG AGGACTTTTCG CAATCACAAG GACCTGTATG ATTTTCGCTA CTTATATCAG	2100
CTCGAGAAAT TGGCATTTAG TACTTATTAT TCATCTTCTG CTTGCTTTCT AAAAAACACT	2160
GCACCTTTAG GTTCTACTCC GCAAGATCCT CTGAGGTTTC AGATGGAAAC ATTGCACTTA	2220
GAGACTCATT CCAGGGCAAC TGCACCTCCA ACTGATGTTT CAACTTTCCT CTTACCTCCT	2280
CCGGATTGTT TTCCACAAAA CCTTAAGAGT TTAACCTTTA GCGGAGATTT CTTTTTGGCA	2340
TGGAAGGATT TGAGCATTGT TGGTAAATTA CCCAACTCG AGGTCCTTCA ACTATCACAC	2400
AATGCCTTCA AAGGCGAGGA GTGGGAAGTA GTTGAGGAAG GGTTTCCTCA CTTGAAGTTC	2460
TTGTTTCTGG ATAGCATATA CATTGCGTAC TGGAGAGCTA GTAGTGATCA CTTTCCATAC	2520
CTTGAACGAC TTTTCTTAG CGATTGCTTT TATTTGGATT CAATCCCTCG AGATTTTGCA	2580
GATATAACCA CACTAGCTCT TATTGATATA TTTCGCTGCC AACAATCTGT TGGGAATTCC	2640
GCCAAGCAAA TTCAACAGGA CATTCAAGAC AACTATGGAA GCTCTATCGA GGTCCATACT	2700
CGTTATCTTT <b>AGTAAGACAT CTTCTTCCTT GATTTACAAC AATATTTAAC TCATCATCAT</b>	2760
<b>AGTAAACTCG ATAATAATCT GGATAATAGC TTTAGTAAGT CAAATTGCAC CAATTCAACA</b>	2820
<b>AAAGTTCTTG ATGCTGTCAT TGTGATTGAT TCGAATCCTT CCAATATTGT GTAACCTGTT</b>	2880
<b>ATACTTGCAT GTTCATTCTT GATTTTGGGA AGTGTAACAT TTCCATTTTT CATCTTGATT</b>	2940
<b>TTGGGAAGTC</b> GAAATGGAGC ATTTTGGTA GTGTGA	2976

10/16

**Fig. 3c (1)**

CTAGAGATTG GAATGGAGTG ATTCTTAGGG GTTTCCTTTTT GAATTAATAT GAGGGTTAGT	60
ATTCAATCTT CAATTCGACA TTTTCTCATA ATTTCTTTAT CTGTTTATTT TTCCTATTCG	120
TAAATCTCTT GGGAAAAATT GGGGTTTTAT CGATTTGGAC TCCTTTTTGA TGAAAAAGGT	180
ATATTTACGA TCTTTATGTT ATGGGTAAAC TGATTTTAAAC ATAAAATTAT TGATTCATCG	240
ATTATTTTTA TCATATTAAC CGCGTACAAT TTGGACTTTC CCGGTAAAGT TAAAGTATGA	300
TAAATTGAGA ATTTCAAGGT CGATCTTAGC TCCATTTTTG ATGAAATTC ATATTTGAAC	360
TTATCTAAGC ATGGGTAAGA TGTTTTTCAA GAAATATTC ATTTTCGAGT CGGGGTTTTG	420
GATTCGAATA TTTTAGGCTT CTTCAAGAAT GTAGATTTTT GTTTAAATTG AGTTTGTGAA	480
TTGATTTCAA CTCCATTTTC AAATTGGTTT TCACCATTAG CTTCCAAATA CTTTAAGGAT	540
CATTTTACAT CAAAAAATTC CAGATTGGG TATCGTTTTC CGGTATGAGA CTTTTGGACC	600
GTTTTGCCCC TTTTCCCTAA ATTTCTTGAT TTTGGTGTCA TTGGACTCGA ATTGTGATTG	660
TGAATAATTG TTTGAATAGA TTATCGTGAT CCAGATTATA CTTGGAAAGG AAAGGCTCAA	720
GTCAAGTAAC TTTTGGAGTT CGTTTTAAGG CAAGTGGCTT CCAAACTTTG TAAAACTCTT	780
AGACTACGCA TGACTACTTT CCTAATTATG TTGGGGAGTA ATGGGGGATT GAGGATGGGT	840
TTTATTTGTT GATTGAAATT GTTGTAATG AAAGATGGGG AATAAAACGA GCTAAATGTG	900
TTATGTGTGA CTTGAATTTG TTTGAATAAG TCATGTGATA ACTGATATTG AGGGATAGAA	960
GAGCATGAGC AGGCTATGAT TGATACAGAC ATTGATGTTG AGGCAGATGA TGTGTAATAC	1020
TATGATGTGG TCGTGATATG GTTGTGATTG AGACATGTGA TGTGTAATAC TATGATGTGG	1080
TCGTGATATG GTTGTGATTG AGACAGGTGA TGTGTAATAC TATGATGTGG TCGTGATATG	1140
GTTGTGATTG AGACAGGTGA TGTGTAATAC TATGATGTGG TCGTGATATG GTTGTGACTG	1200
AGACAGGTGA TGTGTAATAC TATGATGTGG TCGTGATATG GTTGTGATTG AGACAGATGA	1260
TGTGTAATAC GATGATGTGA TCGTGATATG ATTGTGATTG ATTACATGTG CATATTCATT	1320
ATTCATCCCA TGTGTGAAC ATCTGTTGCA TGAGTTCTGA GAACTGATA TGAGGATGGA	1380
TGGATATGAG ACACAGTTGA GACTAGCTCC GGCTAGAGAT GTATGAGATG GACTAGCTCC	1440
GGCTAGCGAT TTGGATGCCG ATGGGATCTG GTTCCGGCGG TGATACATGG TCCATGTGTG	1500
GCCCCCATGG GTTCTGATTT GAGTATTCAA CGCGGACTGA TTACGTCAAC AGATGTGTAT	1560
CGTAGGACAG ACATGTATCA CGACTACATG ACATCATTAT TGCATTTTGC ATCGCATTTG	1620

11/16

**Fig. 3c (2)**

CCTTATCTTT	GTCTGTGATG	TGTGGATTGT	ATCGGTTTAC	CCTTTTTATG	TGGAATTTGA	1680
TCTACTTGCT	CTTATTTGTT	GATCTGAGGT	TGATGAGGAT	ATACTGTTGG	TTCTGGCTGT	1740
TGAATATGAT	CTGTTTAGTA	TAGGTTGGTT	GGTTTGCTGC	TAGATTGAAG	TTTCGGTGGT	1800
TCGGTTGGGA	TTGAAAGGAG	TTGTTTGTAG	CTGCTAGTTT	TGCTTAGTTT	AGAGTTACTT	1860
GCGAGTACCT	GTGGTTTTTCG	GTA CTCACCC	TTGCTTCTAC	ACAATTGTGT	AGGTTGACAG	1920
CTCTCTCTCA	GATATTTTCT	TTAGCAGATT	GAGCTTTGAG	ACATACTCGA	GAGGTAGCGG	1980
TTCATTCCAG	ACGTGCCCTT	GAGTTATCTT	TACTTTTCA GT	TTTGTTCTAT	TCGAGAACTA	2040
TACTCTGAGA	CTTGATATTT	TTTATTCGAA	TTCTGTATTT	AGAGGTTTGT	ACATGTGACA	2100
ACCAAATTCT	GGGTAGTGTT	AAGTCTTAAT	TAAAGTTTTT	TGCTTATTTA	TTATCTTTTA	2160
TTCTCGTATT	TCTACTTCTC	TATCGTTGTG	GTTGGGTTAG	GCTGACGTGT	CTGGTGGGAA	2220
ACGGACATGT	GCCATCACAT	CCGGATTTGG	GGTGTGACAA	ATATTTTGTT	AGTTATATAC	2280
AAAATTGTAT	GTAGTATATG	TATATTTTCT	GCTTTCATCA	CAATTGTATA	TAGATATTTG	2340
TATATTTTGT	TAGTTATATA	CAAAATTGCT	TGAAGTATAT	GTATATTTTC	TGCTTAAATC	2400
ATAATTGTAT	ATATATATAT	ATATATATAT	ATTTCTATAT	TTTGTAAGTT	ATATACAATA	2460
GTATGAATTA	AACAATATAC	AAACCTTACA	TTATTATATA	TACAGTTAGG	TTACACCAAA	2520
AATTATCAAA	TTAAAGCACA	ACTTTTTTTAT	CGAATCATAT	ACAATTCATA	TATATAATTG	2580
ACTTAGTAAT	TTTATACAAC	TACTTACACT	TCTACATGGT	ATAAGAATTT	TGCACAATTA	2640
CTTACATATA	TACAATATTA	TCAATTAAAC	AATATACAAA	TCGTATAACT	TATATATACA	2700
GTAAAATTAC	AACAACAACA	ACAAAAATTA	TCAAATTAAA	GCACACCGTT	GTTGTCGAAT	2760
CATATACACT	CCATATATAC	AAATTGTGTC	ATTCAATTTT	TCGAACAAAA	AATTAGAATT	2820
GAATTGTTAA	TATAAAATTT	ATCTAATATT	GTATAAACAA	AATTAAATTA	TTGCAAACCA	2880
TTAGAATGAA	AAAAACAAAA	ATAAACCGTT	TTCCAAAATT	TCAATTATAT	ACTATACAAA	2940
TCAATTGTAT	ACTTTCTTGC	CGTTCAAAAC	ATGAAGTTTC	CTTGAAAGAA	ACGCTTACCT	3000
AGCGTTGAAT	ATACAAGAAT	ATTGATTAAT	CGTATGCTTC	AGTCGTTTGA	GGAACCCAGT	3060
TGTTATTGTG	TTTCTATTGC	TATAGAACTC	CTTTTTGGAA	AAATATTTGA	TTTGGACGA	3120
TTAGCTTGAA	TCATGGGATT	ATATAAAATT	TTTATTACCG	TATTTAGCAC	TCATGTATCC	3180
ATTTATTAAA	AAAAAATTGT	ATAAATTATA	TTTTTAAAAG	AAAATATACA	AAATTAATGC	3240

12/16

**Fig. 3c (3)**

TTCATAGCAA	ACTAAACTAT	ACCCATTGAA	TGTAATTACT	AAACTATAACC	TATAGAGCGT	3300
TATTTTCATTA	AATACGTTTA	TCATATATGA	AGTTTTCCCT	CAAGAGATCC	TACACCTTAT	3360
ATATAGCTTC	TCAAATGTGG	AAATTCAATC	TCACACCCAA	CAATCTTTCC	CTCAGACTAA	3420
GTTTCATGGC	CCAATATCAC	AATGATCCAC	GAGTCAATTC	ATGAGATTCA	CTATGTGTGT	3480
CACCCACATC	GTCTAAGTAT	TTTATGGCAA	TCAAGCCCTA	CAACTTGCTT	CTTCTTTATA	3540
TATATATATA	TATATATATA	TATATATATA	TATATATGTG	TGTGTGTGTG	TGTGTGTGTG	3600
CGCATCTCTA	ATTAATCTCG	TAAAGGGATT	AAGGGGCCAA	TTTCAAAGAA	TTAGGCGATT	3660
TTCTTAGTTT	TTCGTGTGTG	TTAACCATA	GGTATTTTGG	TGATATGGTT	TTCGGATGAT	3720
TTATTTTGTG	CAACTTATAT	GGAACCCCTC	GTAGGGAGTT	AGTCTCACAC	TTTTTAGAGT	3780
CCATTTTGGG	CATTCAGGGG	CTAATTTATA	GGAAATAGGT	GATCTTCTCA	GTTTGTCTGT	3840
ATTAGCCCAT	GAATATTTTG	GTGATATGTC	TTCCGAATAA	TTTCTTTGTA	AAATCTTTAC	3900
GGGACCCTCC	ATAGGGAGTT	AGTGGAGCAG	TACGTATAGT	CTCACAATTT	TAGAGTTCAT	3960
TTTGGGCATT	TAGGGGCCAA	TTTACAGGAT	TTAGGCGACT	TTCTCAGTGT	TTTGTGTGTG	4020
TTAGCCCATT	AATAGTTGGT	GATATGACTT	TCAGACGATT	TCTTTGCTAC	ACATTTACGG	4080
AACCCTCTGT	AGGAAGTCGG	GGGAGCAATA	CGTACAATCT	CACAATTTTA	GAGTCCATTT	4140
TAGGCATTTA	GGGGCCAATT	TAAAGAAATT	GGACAATTTT	CTCAGTTTTT	CGTGTCTGTT	4200
AGCCATTAAT	ATATTGGTGA	ATATGACCTA	CAGATGATTT	CTAATCGAAA	TCTTTACGAA	4260
ACCCTCAGTA	GGGAGTTGGG	GGAGCAATAC	GTACCGTCTG	ACAATTTTTTA	GAGTCCATTT	4320
TGGGCATTTA	AGGGCCAATT	TACAGGAATT	AGACGATTTT	CTTAGTATTT	TTTCATGTGT	4380
TAGCCCATAA	ATATTTTGTT	GATTTGACTT	TTAGAGTCTA	AACTTCTCAT	GTATATTAAG	4440
AGATATTTAT	GCTTGGTTAA	TTGAATCGAA	CTAGGAATAG	AGAAATTCCT	ACTTGGATCT	4500
TAATATTTCT	CTCTCTTTGA	TTTGGAAAAT	TCTAGGAAGT	TGCTTTCAAT	GGAATTAAAA	4560
TCATCAATCT	CTTGTATGTA	AGAAACATAC	TTATATTCAT	GAATAGATAT	GTTTAGGGTC	4620
TAATAATGAA	TTATCACAAT	TTTTTCTACT	TTTTCTTGTC	AGAGTCCTGC	CTTTTTCTTT	4680
TTCTTTTTTA	ACTTTGGTCT	CTGCTTTTGT	CTACATGATG	ATAAGGTTGG	TGGACCTAGC	4740
TGGAAATGTG	ATGGAAATAG	CTAGTAAAAG	AAAGAACTTT	GCATTTTCTG	TTTTCTTAAA	4800
AACTGATAAA	TTACATAACT	TGTGGCAATT	TGTCCATTTT	CATACTGAGA	GATATTTCTA	4860



13/16

Fig. 3c (4)

TTTTTTTTTG	ATAT <u>ATG</u> GCT	TATGCTGCTG	TTACTTCCCT	TATGAGAACC	ATACATCAAT	4920
CAATGGAACT	TACTGGATGT	GATTTGCAAC	CGTTTTATGA	AAAGCTCAAA	TCTTTGAGAG	4980
CTATTCTGGA	GAAATCCTGC	AATATAATGG	GCGATCATGA	GGGGTTAACA	ATCTTGGAAG	5040
TTGAAATCAT	AGAGGTAGCA	TACACAACAG	AAGATATGGT	TGACTCGGAA	TCAAGAAATG	5100
TTTTTTTTAGC	ACGGAATGTG	GGGAAAAGAA	GCAGGGCTAT	GTGGGGGATT	TTTTTCGTCT	5160
TGGAACAAGC	ACTAGAATGC	ATTGATTCCA	CCGTGAAACA	GTGGATGGCA	ACATCGGACA	5220
GCATGAAAGA	TCTAAAACCA	CAAAC TAGCT	CAC TTGTCAG	TTTACCTGAA	CATGATGTTG	5280
AGCAGCCCGA	GAATATAATG	GTTGGCCGTG	AAAATGAATT	TGAGATGATG	CTGGATCAAC	5340
TTGCTAGAGG	AGGAAGGGAA	CTAGAAGTTG	TCTCAATCGT	AGGGATGGGA	GGCATCGGGA	5400
AAACAAC TTT	GGCTGCAAAA	CTCTATAGTG	ATCCTTACAT	TATGTCTCGA	TTTGATATTC	5460
GTGCAAAAGC	AACTGTTTCA	CAAGAGTATT	GTGTGAGAAA	TGTACTCCTA	GGCCTTCTTT	5520
CTTTGACAAG	TGATGAACCT	GATTATCAGC	TAGCGGACCA	ACTGCAAAAG	CATCTGAAAG	5580
GCAGGAGATA	CTTG GTAGTC	ATTGATGACA	TATGGACTAC	AGAAGCTTGG	GATGATATAA	5640
AACTATGTTT	CCCAGACTGC	GATAATGGAA	GCAGAATACT	CCTGACTACT	CGGAATGTGG	5700
AAGTGGCTGA	ATATGCTAGC	TCAGGTAAGC	CTCCTCATCA	CATGCGCCTC	ATGAATTTTG	5760
ACGAAAGTTG	GAATTTACTA	CACAAAAGAA	TCTTTGAAAA	AGAAGGTTCT	TATTCTCCTG	5820
AATTTGAAAA	TATTGGGAAA	CAAATTGCAT	TAAAATGTGG	AGGGTTACCT	CTAGCAATTA	5880
CTTTGATTGC	TGGACTTCTC	TCCAAAATCA	GTAAACATT	GGATGAGTGG	CAAAATGTTG	5940
CGGAGAATGT	ACGTTCCGGT	GTAAGCACAG	ATCTTGAAGC	AAAATGCATG	AGAGTGTTGG	6000
CTTTGAGTTA	CCATCACTTG	CCTTCTCACC	TAAAACCGTG	TTTTCTGTAT	TTTGCAATTT	6060
TCGCAGAGGA	TGAACGGATT	TATGTAAATA	AACTTGTTGA	GTTATGGGCC	G TAGAGGGGT	6120
TTTTGAATGA	AGAAGAGGGA	AAAAGCATAG	AAGAGGTGGC	AGAAACATGT	ATAAACGAAC	6180
TTGTAGATAG	AAGTCTAATT	TCTATCCACA	ATGTGAGTTT	TGATGGGGAA	ACACAGAGAT	6240
GTGGAATGCA	TGATGTGACC	CGTGAACTCT	GTTTGAGGGA	AGCTCGAAAC	ATGAATTTTG	6300
TGAATGTTAT	CAGAGGAAAG	AGTGATCAAA	ATTCATGTGC	ACAATCCATG	CAGTGTTCCCT	6360
TTAAGAGTCG	AAGTCGGATC	AGTATCCATA	ATGAGGAAGA	ATTGGTTTGG	TGTCGTAACA	6420
GCGAGGCTCA	TTCTATCATC	ACGTTGTGTA	TATTCAAATG	CGTCACACTG	GAATTGTCTT	6480

14/16

Fig. 3c (5)

TCAAGCTAGT AAGAGTACTA GATCTTGGTT TGACTACATG CCCAATTTTT CCCAGTGGAG	6540
TACTTTCTCT AATTCATTTG AGATACCTAT CTTTGCGTTT TAATCCTCGC TTACAGCAGT	6600
ATCGAGGATC GAAAGAAGCT GTTCCCTCAT CAATAATAGA CATTCTCTA TCGATATCAA	6660
GCCTATGCTA TCTGCAAAC TTTAAACTTT ACCATCCATT TCCCAATTGT TATCCTTTCA	6720
TATTACCATC GGAAATTTTG ACAATGCCAC AATTGAGGAA GCTGTGTATG GGCTGGAATT	6780
ACTTGCGGAG TCATGAGCCT ACAGAGAACA GATTGGTTTT GAAAAGTTTG CAATGCCTCA	6840
ATGAATTGAA TCCTCGGTAT TGTACAGGGT CTTTTTTAAG ACTATTTCCC AATTAAAGA	6900
AGTTGGAAGT ATTTGGCGTC AAAGAGGACT TTCGCAATCA CAAGGACCTG TATGATTTTC	6960
GCTACTTATA TCAGCTCGAG AAATTGGCAT TTAGTACTTA TTATTCATCT TCTGCTTGCT	7020
TTCTAAAAA CACTGCACCT TTAGGTTCTA CTCCGCAAGA TCCTCTGAGG TTTCAGATGG	7080
AAACATTGCA CTTAGAGACT CATTCCAGGG CAACTGCACC TCCAACGAT GTTCCAACCT	7140
TCCTCTTACC TCCTCCGGAT TGTTTTCCAC AAAACCTTAA GAGTTTAACT TTTAGCGGAG	7200
ATTTCTTTTT GGCATGGAAG GATTTGAGCA TTGTTGGTAA ATTACCCAAA CTCGAGGTCC	7260
TTCAACTATC ACACAATGCC TTCAAAGGCG AGGAGTGGGA AGTAGTTGAG GAAGGGTTTC	7320
CTCACTTGAA GTTCTTGTTT CTGGATAGCA TATACATTCG GTACTGGAGA GCTAGTAGTG	7380
ATCACTTTCC ATACCTTGAA CGACTTTTTC TTAGCGATTG CTTTATTTG GATTCAATCC	7440
CTCGAGATTT TGCAGATATA ACCACACTAG CTCTTATTGA TATATTTTCG TGCCAACAAT	7500
CTGTTGGGAA TTCCGCCAAG CAAATTCAAC AGGACATTCA AGACAACTAT GGAAGCTCTA	7560
TCGAGGTCCA TACTCGTTAT CTTAGTAAG ACATCTTCTT CCTTGATTTA CAACAATATT	7620
TAACTCATCA TCATAGTAAA CTCGATAATA ATCTGGATAA TAGCTTTAGT AAGTCAAATT	7680
GCACCAATTC AACAAAAGTT CTTGATGCTG TCATTGTGAT TGATTCTGAAT CCTTCCAATA	7740
TTGTGTAAC TGTTATACTT GCATGTTTCAT TCTTGATTTT GGGAAAGTGTA ACATTTCCAT	7800
TTTTCATCTT GATTTTGGGA AGTCGAAATG GAGCATTTTT GGTAGTGGA CAACAGATGA	7860
AGATGATGAT GATAGTGTGA CAACAGATGA AGATGAAGAT GAAGACTTTG AGAAAGAAGT	7920
TGCTTCTTGC GGCAATAATG TGTAAGTTCT TATACCTGCA TGCTCATTCT TGCTATAATG	7980
TTCTCTTGTT CCTTAATTAT GGGACATCTA ACATATTATT TTCCATTTTT TGCATCTTTT	8040
TTTTTTCCTG CAGCGTGTAG TTAAGGTGTT CTGAGGACTA GCCAGTTCTC TGAAATAAAT	8100

15/16

**Fig. 3c (6)**

GTCAAATCAG	AAGCCAAATG	TGTGAGTGTT	TGTTTTGTTC	GTTTTCATTT	TTTCTGCATA	8160
AGGTGGCAGG	ATGATTGCAA	ATGGCTTGTA	ATTTAATTGT	ATATGATATT	TCGTATAGCC	8220
ATTTGCCAGT	GGTTTTTTTAG	ATACTCCAAA	TTTTATGTAC	ATACATAATG	GTATAGGCCA	8280
GAACAGGCTC	CATATATAAC	GTGTGTTTCC	TTTCTTGGA	GTCCTCAATC	TACCTCGCAA	8340
AGGAAGACAG	ACGGCTAAAT	CAAGAAAGAA	ATTTTTTTGA	AAATCATGTG	GCTAGTTGTT	8400
CAACTTTATA	CAAGTTTATG	TGCATACTTG	TGCATACCCA	AAGTTGAATA	ACATAAACAT	8460
AAAATGAAGT	CAAGTTAAAT	GGCACATTTA	TGTATTATGC	CTTTTGAATT	TCATTAATAG	8520
TGAAAATCCT	GAATCATATT	CAGATTCCAT	CACTAATCGT	TGAACCATGT	TAATTTACTA	8580
TGTATTATCT	AATGGATTTT	TTTGCTATCT	TATTTATAAT	TGTTCAAAGT	TTTGTTAATT	8640
ATCTTTAGCA	TAATATCTGA	TTATATTATT	TTGATATACT	TTCTCTATCC	CTAATTACTT	8700
GTCCATTTTT	GAATTGGCAC	ACCTATTAAG	AAAATAATTA	TTGAAATAGT	GAGTTTACCA	8760
TTTTACCCAT	ATTAATTATG	AAGTGGATGA	ATTAAAAACT	CAAGATTTTC	AAAAAGTTCT	8820
ATTTTTTTCA	AAGTAATAAA	CTGACGGTAT	AATAGGTAAA	AAAAATTATT	CTTTCTTGAT	8880
TTGTCAAAT	AAACAAATAA	TTAGGAATAA	TTAAAAAAT	GGATAAATAA	TTAAAAACGG	8940
AGGGAGCAAT	ATGTTATCTT	TAGCCTAATA	ATATCTGATT	AATGGCCACC	CTAATTGATT	9000
GGATAGGAGA	GGATAGACTT	GCTTCCAAGT	AACCCAAAAT	ATAAAAAGTT	GACAAAAGGG	9060
TGCTAAATTC	GAGACACATG	TAGTACTTAT	ATAATTCATG	TGCGGACTCG	TTCTTTTGTA	9120
GTACTCCCTC	CGTTCTATTT	TATACGTCAC	ATTTTTACTT	TATACTTTTA	TTAAGAAATG	9180
ATGTAGTTTT	ATCTTTCTAT	TCTTATTTAA	TGTTTTCTTA	AGTCAATTTT	ATAATAAATA	9240
ATGAATATAT	TTTCAAGATT	AATTAACTAC	TCTATCAAGG	GTATAATAGG	TAAAATATGA	9300
TAATTTATAC	ATAAATTTTA	TAAAATGACA	AGTATTGTGG	TCCAACCTATT	TATAGAAAGA	9360
AATGATATAT	AAAATGGGAC	GGAGGGCGTT	ATAAAGTTGA	CTTAAGAAAA	CATTAAATAA	9420
GGGTAGAAGG	GTAAAATTAC	ATTATTTCTT	AATGTAAATG	TAAAGTAAAA	AGGTAACATA	9480
TAAAATGGAA	AGGAGGGAGT	AGTATTTTCT	TGTTTTATTT	TACGTGGCAC	TCTATTCTCA	9540
TAATCCGTCT	TTAAAAATGT	CATTTTATTG	TAATTGAAAA	TAATTTAACT	TAAAATTCTC	9600
CATCTACCCT	TAATTAATGA	AATGATTTAC	AATTATATAA	ATATATAAAA	ATTGTTTTAG	9660
CCTATAATTT	TCTAAAATCT	TTTTTTTTCT	CTTATACATC	GTATTAAGTC	AAACATAAAT	9720

16/16

**Fig. 3c (7)**

GGAATGGACG	GAGTATTTCT	TTTATTTTTT	TGTCACACCG	CCCATATGTT	TTCTCCCATC	9780
CCCCAGACCC	CCACTATGTA	TATTTACTCC	TTAGTTGGAT	CTGAATTTAG	AGTTTAGAAG	9840
CTTCTATAAT	AATTTTAGAT	TAATATATAA	TAATAATAAT	AATAATTGAA	CTTACAGTAT	9900
TAAATTTATG	TGAATCTATA	TATATTGTAT	TGTAATTTTT	TTAATTATAA	TTTTAACCAA	9960
ATCAATAAAG	CTATTCAGAT	GTAAAAGTAT	ATATTATGAT	TTAACAACAA	ATTTCTATAC	10020
GTCTTCCTAA	GTTTTGATGC	ATAATTTCTT	AAAACATATA	AATTTCCAAG	TGACTACTTC	10080
CAGTATTACA	ATGAGAACTT	ATGTTTCGTT	ATGGATTTTC	TTAGTGAATT	AGTTTAATAA	10140
AATCAAAATG	AAAAAAAATC	ATGTTTTATA	ACATAAAATT	TTCATTGATT	CATGCGAAAA	10200
AAAAACATCT	AGTTCTTATA	GTGTGAAAAC	TATTGAACTT	ATGGGATGTA	GCTGTATGGA	10260
AGTTCATCAA	GTGGTAGCTC	CTTGTACGCA	ACTAGTGCTA	CTTTTATTG	ACTAAAAGTT	10320
ATTTTCTAG						10329